arXiv:math/0002062v1 [math.CO] 9 Feb 2000

A characterisation of Pfaffian near bipartite graphs

Ilse Fischer
Universität Klagenfurt
A-9020 Klagenfurt
Austria
Email: `Ilse.Fischer@uni-klu.ac.at`

C.H.C. Little[1]
Massey University
Palmerston North
New Zealand
Email: `c.little@massey.ac.nz`

---

[1] The second author thanks the University of Klagenfurt for its hospitality while this research was undertaken.

PFAFFIAN NEAR BIPARTITE GRAPHS

Ilse Fischer
Universität Klagenfurt
Universitätsstrasse 65-67
A-9020 Klagenfurt
Austria
Email: `Ilse.Fischer@uni-klu.ac.at`

Abstract

A graph is *1-extendible* if every edge has a 1-factor containing it. A 1-extendible non-bipartite graph G is said to be *near bipartite* if there exist edges e_1 and e_2 such that $G - \{e_1, e_2\}$ is 1-extendible and bipartite. We characterise the Pfaffian near bipartite graphs in terms of forbidden subgraphs. The theorem extends an earlier characterisation of Pfaffian bipartite graphs.

1. Introduction

The graphs considered in this paper are finite and have no loops or multiple edges. They are also undirected and connected unless an indication to the contrary is given. If v and w are vertices in a directed graph, then (v, w) denotes an edge joining v and w and directed from v to w. If G is any graph, then we denote its vertex set by VG and its edge set by EG. A 1-*factor* of G is a subset f of EG such that every vertex has a unique edge of f incident on it.

Let G^* be a directed graph with an even number $2n$ of vertices and let F be the set $\{f_1, f_2, \ldots, f_k\}$ of 1-factors of G^*. For all i write

$$f_i = \{(u_{i1}, w_{i1}), (u_{i2}, w_{i2}), \ldots, (u_{in}, w_{in})\},$$

where $u_{ij}, w_{ij} \in VG^*$ for all j. Associate with f_i a plus sign if

$$u_{i1}w_{i1}u_{i2}w_{i2} \ldots u_{in}w_{in}$$

is an even permutation of

$$u_{11}w_{11}u_{12}w_{12} \ldots u_{1n}w_{1n},$$

and a minus sign otherwise. Note that the signs of the 1-factors are independent of the order in which their edges have been written. They are dependent on the choice of f_1, but the resulting partition of F into two complementary subsets is not. If G is an undirected graph, we say that G is a *Pfaffian* graph if there exists an orientation such that all the 1-factors of G have the same sign. We say that this orientation is a *Pfaffian* orientation of G. Pfaffian orientations have been used by Kasteleyn [1] to enumerate 1-factors in planar graphs. In fact his method can be used precisely for those graphs that are Pfaffian. It is therefore of interest to know which graphs are Pfaffian, but this question is open.

Pfaffian bipartite graphs have been characterised by Little [3], who proved the following theorem.

Theorem 1. *A bipartite graph G is non-Pfaffian if and only if it contains an even subdivision J of $K_{3,3}$ such that $G - VJ$ has a 1-factor.*

Here we need to explain the term ‘even subdivision’. An *edge subdivision* of a graph G is defined as a graph obtained from G by replacing an edge joining vertices v and w with a path P joining v and w but having no other vertices in common with G. The edge subdivision is *even* if P has odd length. A graph H is a *subdivision* of G if for some positive integer k there exist graphs $G_0, G_1, \ldots, G_k$ such that $G_0 = G$, $G_k = H$ and, for all $i > 0$, G_i is an edge subdivision of G_{i-1}. If $G_1, G_2, \ldots, G_k$ can be chosen so that in addition G_i is an even edge subdivision of G_{i-1} for all $i > 0$, then H is said to be an *even subdivision* of G. It is easy to see that G is Pfaffian if and only if H is Pfaffian. A more general result is proved in Lemma 2.

At this point it is worth mentioning that Robertson, Seymour and Thomas [6] have recently found a polynomial-time algorithm which decides whether a bipartite graph is Pfaffian or not.

A graph is *1-extendible* if every edge has a 1-factor containing it. Such graphs are the only graphs of interest in the study of the Pfaffian property, as any edge belonging to no 1-factor is irrelevant. A 1-extendible non-bipartite graph G is said to be *near* *bipartite* if there exist edges e_1, e_2 such that $G - \{e_1, e_2\}$ is 1-extendible and bipartite. If G were a 1-extendible graph and $G - \{e\}$ were bipartite for some edge e, then G would also be bipartite. This observation explains why we remove two edges from G, rather than one, in the definition of a near bipartite graph. The aim of the present paper is to extend Theorem 1 to a characterisation of Pfaffian near bipartite graphs in terms of forbidden subgraphs.

In the statement of our main theorem below, Γ_1 and Γ_2 refer to the graphs drawn in Figures 2 and 3 respectively, where the arrows are to be ignored. Both graphs are near bipartite, since $\Gamma_1 - \{(f,l),(i,c)\}$ and $\Gamma_2 - \{(f,e),(i,j)\}$ are 1-extendible and bipartite. Note that Γ_2 may be obtained from the Petersen graph by subdividing two fixed edges at a maximum distance apart and then joining the vertices of degree 2 by an edge. These graphs, like $K_{3,3}$, can easily be shown to be non-Pfaffian. Indeed, each graph in Figures 1–3 is accompanied by a set S of 1-factors such that each edge belongs to just two members of S and S contains an odd number of 1-factors of each kind of sign under the given orientation. The former property of S implies that the latter is still valid if we change the orientation of a single edge. Therefore the latter property of S is independent of the orientation and consequently the graphs cannot be Pfaffian.

It follows that no even subdivision of these graphs is Pfaffian. It is shown in [4] that a graph G is non-Pfaffian if it has a circuit X, of odd length, such that the graph obtained from G by contracting X to a vertex is non-Pfaffian. In general, let us say that a graph G is *simply reducible* to a graph H if G has a circuit X, of odd length, such that H is obtained from G by contracting X. More generally, we say that G is *reducible* to a graph H if for some positive integer k there exist graphs $G_0, G_1, \dots, G_k$ such that $G_0 = G$, $G_k = H$ and, for all $i > 0$, G_{i-1} is simply reducible to G_i. Thus any graph that is reducible to an even subdivision of $K_{3,3}$, Γ_1 or Γ_2 is non-Pfaffian. In fact, a graph G must be non-Pfaffian if it has a subgraph J that is reducible to an even subdivision of $K_{3,3}$, Γ_1 or Γ_2 and has the property that $G - VJ$ has a 1-factor. The purpose of this paper is to show that the converse of this statement holds for near bipartite graphs.

Theorem 2. *A near bipartite graph G is non-Pfaffian if and only if G contains a subgraph J, reducible to an even subdivision of $K_{3,3}$, Γ_1 or Γ_2, such that $G - VJ$ has a 1-factor.*

Definitions and Notation. The following definitions and notation are fundamental for this paper. Circuits, non-empty paths and, more generally, subgraphs with no isolated vertices are determined by their edge sets, and are therefore identified with them in this paper. If X is a path or circuit in a graph G, then we denote by VX the set of vertices of X. If P is a path and $u, v \in VP$, then we denote by $P[u,v]$ the subpath of P joining u to v. If $P[u,v]$ is directed from u to v, then we also write $P(u,v) = P[u,v]$. If C is a circuit which includes a unique directed path from vertex u to vertex v, then that path is denoted by $C(u,v)$. From time to time we may perform a reorientation of C, that is to say we change the orientation of every oriented edge in C. This directed path is then denoted by $C(v,u)$, or by $P(v,u)$ if it is included in another path P.



$(a,b)(c,d)(e,f)$ $+$
$(a,b)(c,f)(e,d)$ $-$
$(a,d)(c,f)(e,b)$ $+$
$(a,d)(c,b)(e,f)$ $-$
$(a,f)(c,b)(e,d)$ $+$
$(a,f)(c,d)(e,b)$ $-$

FIGURE 1. The graph $K_{3,3}$.



$(a,b)(c,d)(e,f)(g,h)(i,j)(k,l)$ $+$
$(b,c)(d,e)(f,g)(h,i)(j,k)(l,a)$ $-$
$(d,a)(j,e)(b,c)(f,g)(h,i)(k,l)$ $+$
$(b,g)(h,k)(c,d)(e,f)(i,j)(l,a)$ $-$
$(i,c)(f,l)(a,b)(d,e)(g,h)(j,k)$ $+$
$(d,a)(b,g)(i,c)(j,e)(h,k)(f,l)$ $-$

FIGURE 2. The graph Γ_1.

A circuit is *alternating* with respect to each of two given 1-factors if it is included in their symmetric difference. A circuit that is alternating with respect to a 1-factor f is also said to be f*-alternating*, or *consanguineous* (with respect to f). Note that a graph with more than one edge is 1-extendible if and only if every edge has an alternating circuit containing it. A path P is *alternating* if every internal vertex of P is incident with an edge of $P \cap f$. An *ear* is a path of odd cardinality.

Let A and B be sets of edges in a graph G. Then an AB*-arc* is a non-empty maximal subpath of $A \cap B$, and an $A\overline{B}$*-arc* (or a $\overline{B}A$*-arc*) is a non-empty maximal subpath of $A - B$. A $G\overline{B}$-arc is also called a $\overline{B}$-arc.



$(a,b)(c,d)(f,e)(h,g)(i,j)(k,l)$	$+$
$(b,c)(d,e)(g,f)(i,h)(j,k)(l,a)$	$-$
$(e,l)(a,b)(c,d)(g,f)(i,h)(j,k)$	$+$
$(h,d)(k,g)(b,c)(f,e)(i,j)(l,a)$	$-$
$(a,i)(f,b)(c,j)(d,e)(h,g)(k,l)$	$+$
$(a,i)(f,b)(c,j)(h,d)(e,l)(k,g)$	$-$

FIGURE 3. The graph Γ_2.

2-Ear Theorem. Next, let A be an alternating circuit in G and let H be a subgraph of G. If there are n $A\overline{H}$-arcs, and each is an ear, then we say that $G[EH \cup A]$ is obtained from H by an *n-ear adjunction.* An *ear decomposition* of a 1-extendible graph G is a sequence $G_0, G_1, \dots, G_t$ of 1-extendible subgraphs of G such that G_0 is isomorphic to K_2, $G_t = G$ and, for each $i > 0$, G_i is obtained from G_{i-1} by a 1-ear or 2-ear adjunction. A theorem of Lovász and Plummer [5, Theorem 5.4.6] asserts that every 1-extendible graph has an ear decomposition. It can be stated as follows.

Theorem 3. *Let f be a 1-factor in a 1-extendible graph G. Let H be a 1-extendible proper subgraph of G such that $EH \neq \emptyset$ and $f \cap EH$ is a 1-factor of H. Then G contains an f-alternating circuit A that admits just one or two $A\overline{H}$-arcs.*

In fact if G is bipartite then it can be shown that only 1-ear adjunctions are necessary.

The idea behind the proof of Theorem 1 runs as follows. Clearly we may assume that G is 1-extendible. Suppose that G is non-Pfaffian. We construct an ear decomposition $G_0, G_1, \dots, G_t$ of G. Since G is bipartite, we may assume that, for each $i > 0$, G_i is obtained from G_{i-1} by the adjunction of a single ear. As G_0 is Pfaffian but G is not, there exists a smallest positive integer j such that G_j is non-Pfaffian. The graph G_j is studied in detail and eventually shown to contain J.

Theorem 3 provides a possible way to generalise this argument. If we drop the assumption that G is bipartite then, for each i, G_i is obtained from G_{i-1} by the adjunction of one or two ears. In this paper we consider the case where G_{j-1} is bipartite and G_j is obtained from G_{j-1} by a 2-ear adjunction.

Idea behind the proof of Theorem 2. We use alternating circuits in preference to 1-factors. Kasteleyn [1] has shown that the 1-factors of a directed graph all have equal sign if and only if all the alternating circuits are clockwise odd. (The clockwise parity of a circuit of even length is the parity of the number of its edges that are directed in agreement with a specified sense.) Let G be a near bipartite graph which is minimal with respect to the property of being non-Pfaffian. Let e_1 and e_2 be edges of G such that $G - \{e_1, e_2\}$ is bipartite and 1-extendible. By minimality $G - \{e_1, e_2\}$ has a Pfaffian orientation. Extend this orientation to an orientation of G by orienting e_1 and e_2 arbitrarily. Since G is non-Pfaffian, there exist two alternating circuits A and B of opposite clockwise parity. In Theorem 5 we construct alternating circuits in $G - \{e_1, e_2\}$ whose union is $EG - \{e_1, e_2\}$ and whose sum (symmetric difference) is $A + B$. This construction is used to generate all the non-Pfaffian near bipartite graphs. The list of non-Pfaffian near bipartite graphs so constructed is infinite. In Sections 3 and 4 we are then able to reduce this list to a finite list by invoking the minimality of G. In Section 5 we finally show that every graph in this list can be obtained from $K_{3,3}$, Γ_1 or Γ_2 by means of the operations of reduction and even subdivision. In Section 6 we demonstrate that neither Γ_1 nor Γ_2 is reducible to an even subdivision of $K_{3,3}$.

2. A STRUCTURE THEOREM OF MINIMAL NON-PFAFFIAN NEAR BIPARTITE GRAPHS

In this section we establish that a minimal non-Pfaffian near bipartite graph is the union of two alternating circuits A and B and two additional paths S and T. Let G be a near bipartite graph. We may assume that G is minimal with respect to the property of being non-Pfaffian. To see this point, suppose that G has an edge e such that $G - \{e\}$ is non-Pfaffian and has a subgraph J, reducible to an even subdivision of $K_{3,3}, \Gamma_1$ or Γ_2, such that $(G - \{e\}) - VJ$ has a 1-factor f. Then f is also a 1-factor of $G - VJ$, and so Theorem 2 holds also for G.

A set S of alternating circuits in a directed graph H is called *intractable* if the sum of the circuits in S is empty and an odd number of the members of S are clockwise even. The former property implies that the latter is independent of the orientation of H. (See Lemma 8.) The following lemma is proved in [2].

Lemma 1. *A graph is Pfaffian if and only if it has no intractable set of alternating circuits.*

From this result we show that we can assume there to be no vertices of degree 2 in G.

Lemma 2. *Let v be a vertex of degree 2 in G, and let G' be the graph obtained from G by contracting the edges incident on v. Then G is Pfaffian if and only if G' is Pfaffian.*

Proof: Let a and b be the edges of G incident on v, and let u and w be the vertices adjacent to v.

Suppose there is an intractable set S of alternating circuits in G. Then the intersections of the circuits in S with $EG - \{a, b\}$ yield an intractable set in G'. Conversely, let S' be an intractable set of alternating circuits in G'. Let v' be the vertex in G' obtained by identifying u and w in G. Choose $C' \in S'$. If $v' \notin VC'$, or the edges of C' incident on v' in G' are both incident on u in G or both incident on w in G, then let $C = C'$; otherwise let $C = C' \cup \{a, b\}$. The set S of such circuits C forms an intractable set in G. (Note that the sum of the circuits in S is a subset of $\{a, b\}$ and therefore empty as it must be a cycle.) ☐

Let G be a graph with a vertex of degree 2 and let G' be the graph obtained from G by contracting the edges incident on it. Suppose that in G' there is a subgraph J, reducible to an even subdivision of $K_{3,3}, \Gamma_1$ or Γ_2, such that $G' - VJ$ has a 1-factor.

Then the same is true for G, for $K_{3,3}$, Γ_1 and Γ_2 are cubic and so the converse of the reduction in the lemma gives an even subdivision of each of those graphs. Therefore we can assume that G has no vertex of degree 2.

Since G is near bipartite, it is 1-extendible. Moreover there exist edges e_1 and e_2 such that $G - \{e_1, e_2\}$ is bipartite and 1-extendible. We call this graph H, and fix a 1-factor f of H. Note that $G - \{e_1\}$ is non-bipartite, for otherwise, since G is non-bipartite, every circuit containing e_1 would be of odd length, in contradiction to the fact that G has an alternating circuit containing e_1. Similarly $G - \{e_2\}$ is non-bipartite. Consequently any alternating circuit containing one of e_1 and e_2 must also contain the other.

Note that H is Pfaffian, by the minimality of G. Extend a Pfaffian orientation of H to an orientation of G by orienting e_1 and e_2 arbitrarily. We shall henceforth refer to this orientation as our extended Pfaffian orientation of G. As G is non-Pfaffian, it contains a clockwise even alternating circuit A. This circuit must contain e_1 and e_2. There must also be a clockwise odd alternating circuit B containing e_1 and e_2, for otherwise a Pfaffian orientation for G could be constructed by reorienting e_1 or e_2. The following lemma, which is proved in [4], gives information about how A and B can be chosen.

Lemma 3. *Let f be a 1-factor in a 1-extendible directed graph G. Let A and B be f-alternating circuits in G, of opposite clockwise parity, containing distinct independent edges e_1 and e_2 such that $e_1 \notin f$ and $e_2 \notin f$. Suppose that $G - \{e_1\}$ and $G - \{e_2\}$ are not bipartite but that $G - \{e_1, e_2\}$ is. Then $A \cup B$ includes alternating circuits X and Y, of opposite clockwise parity and consanguineous with respect to some 1-factor that contains neither e_1 nor e_2, such that there are just one or two XY-arcs, each XY-arc contains e_1 or e_2 and their union contains both.*

Thus A and B can be chosen so that there are at most two AB-arcs. In [4] the case where there is a unique AB-arc has been dealt with. We obtained the following theorem.

Theorem 4. *Let G be a 1-extendible graph with 1-factor f. Let e_1 and e_2 be distinct independent edges of $EG - f$ such that neither $G - \{e_1\}$ nor $G - \{e_2\}$ is bipartite but $G - \{e_1, e_2\}$ is bipartite, Pfaffian and 1-extendible. Suppose there exist f-alternating circuits A and B, both containing e_1 and e_2, such that there is a unique AB-arc and A and B have opposite clockwise parity under a Pfaffian orientation of $G - \{e_1, e_2\}$. Then G has a subgraph J, reducible to an even subdivision of $K_{3,3}$, such that $G - VJ$ has a 1-factor.*

In the present paper, we deal with the remaining case, where for every choice of A and B there are at least two AB-arcs. Henceforth we assume that A and B are chosen so that there are exactly two AB-arcs, and therefore exactly two $\overline{A}B$-arcs and exactly two $A\overline{B}$-arcs. By Lemma 3 it may be assumed that one of the AB-arcs contains e_1 and the other e_2. Let the former arc join vertices x_1 and x_2 and the latter vertices y_1 and y_2. Let e_1 join vertices u_1 and u_2 and let e_2 join vertices v_1 and v_2. Define $A^* = A - \{e_1\}$, and adjust the notation so that the vertices u_1, x_1, y_1, v_1, v_2, y_2, x_2, u_2 appear in that order as A^* is traversed from u_1 to u_2.



FIGURE 4. A homeomorph of $G[A \cup B]$.

Lemma 4. *One of the* $\overline{A}B$*-arcs joins* x_2 *to* y_1 *and the other* x_1 *to* y_2.

Proof: Suppose the contrary. Note that the edges of f incident on x_1, x_2, y_1 and y_2, respectively, belong to $A^*[x_1, u_1] \cup A^*[u_2, x_2] \cup A^*[y_2, y_1]$, since e_1 and e_2 belong to AB-arcs. If an $\overline{A}B$-arc X were to join y_1 to y_2 then we should have the contradiction that the circuit $A^*[y_1, y_2] \cup X$ would be of even length yet contain e_2 but not e_1. On the other hand, suppose that an $\overline{A}B$-arc Y were to join x_1 to y_1. Let

$$C = Y \cup A^*[x_1, u_1] \cup \{e_1\} \cup A^*[u_2, y_1].$$

This circuit, an f-alternating circuit containing e_1 and e_2, would have opposite clockwise parity from either A or B. Since there would be a unique AC-arc and a unique BC-arc, we should have a contradiction to the assumption that there is no choice for A and B that gives a unique AB-arc. ☐

The graph $G[A \cup B]$ is an even subdivision of that shown in Figure 4. The edges of f are thickened in this and subsequent figures, and in all subsequent figures the graph in question is an even subdivision of the one portrayed.

For a bipartite graph K with bipartition $\{M, N\}$ and 1-factor f there exists an orientation in which the directed paths and directed circuits are precisely the f-alternating paths and f-alternating circuits respectively: orient the edges of f from M to N and the remaining edges from N to M. Then every vertex has indegree 1 or outdegree 1, and every edge joins a vertex of indegree 1 to a vertex of outdegree 1. It follows that directed paths with an internal vertex in common meet in an edge incident on the vertex. We call this orientation the *reference* orientation for K with respect to (M, N, f). Fix such an orientation for H so that $A^*[u_1, v_1]$ is directed from u_1 to v_1. We refer to this orientation as our reference orientation for H. It follows that $B \cap EH$ includes a directed path from u_1 to v_2 and another from u_2 to v_1, and that $A^*[u_2, v_2]$ is directed from u_2 to v_2. The orientation is also indicated in Figure 4. Henceforth the



FIGURE 5. $A \cup B \cup S \cup T$

orientation associated with H will be our reference orientation unless an indication to the contrary is given.

Let f' be another 1-factor of K. It is shown in [3] that the reference orientation for K with respect to (M, N, f') is obtained from the reference orientation with respect to (M, N, f) by reorienting the circuits included in $f + f'$. This fact is used implicitly later on to justify reorientations of f-alternating circuits.

The following lemma is a standard result. (See [3].)

Lemma 5. *Let G be a directed graph such that each edge has a directed circuit containing it. Then for every $a, b \in VG$, there exists a directed path from a to b.*

We may apply this lemma to H, since every edge of the 1-extendible graph H must belong to a directed circuit. Thus there is a directed path S from y_1 to x_1 and a directed path T from y_2 to x_2. (See Figure 5; a dotted line in this and subsequent figures stands for a directed path, which can have intersections with the rest of the graph that are not indicated.) We now aim to show that $EG = A \cup B \cup S \cup T$ in Theorem 5. To this end we introduce the following three lemmas.

Lemma 6. *Let $(a_1, a_2, \ldots, a_n)$ be a sequence of edges in a directed graph G in which each vertex has indegree 1 or outdegree 1. Suppose that for all $i > 1$ the origin of a_i is the terminus of a_{i-1} and that the origin of a_1 has outdegree 1 and the terminus of a_n has indgree 1. Then there exist a directed path P from the origin of a_1 to the terminus of a_n and directed circuits $C_1, C_2, \ldots, C_k$ such that*

$$\sum_{i=1}^{n}\{a_i\} = P + \sum_{i=1}^{k} C_i$$

and

$$P \cup \bigcup_{i=1}^{k} C_i = \{a_1, a_2, \ldots, a_n\}.$$

Proof: We use induction on the number r of repetitions of edges in the sequence $L = (a_1, a_2, \ldots, a_n)$. If $r = 0$ then $P = \{a_1, a_2, \ldots, a_n\}$ and $k = 0$, since each vertex has indegree 1 or outdegree 1, the origin of a_1 has outdegree 1 and the terminus of a_n has indegree 1. Now suppose that $r > 0$ and that the lemma holds whenever the number of repetitions of edges is less than r. Let a be the edge in L that is repeated first. The part of L between the first two occurrences of a has no edges repeated within it. Therefore a and the edges between the first two occurrences of a form a directed circuit C. We now modify L by removing all the edges from the first occurrence of a to the edge immediately before the second occurrence of a. This modified sequence $L' = (a'_1, a'_2, \ldots, a'_m)$ has fewer repetitions of edges. Moreover the origin of a'_1 is that of a_1, the terminus of a'_m is that of a_n, and for all $i > 1$ the origin of a'_i is the terminus of a'_{i-1}. Therefore the inductive hypothesis may be applied to L', and the result follows from the equation

$$\sum_{i=1}^{n} \{a_i\} = \sum_{i=1}^{m} \{a'_i\} + C.$$

□

Lemma 7. *Let H be a directed graph, let P be a directed path from vertex x to vertex y and let Q be a directed path from y to x. Then there are directed circuits $C_1, C_2, \ldots, C_k$ such that*

$$\sum_{i=1}^{k} C_i = P + Q$$

and

$$\bigcup_{i=1}^{k} C_i = P \cup Q.$$

Proof: We use induction on $n = |VP \cap VQ| \geq 2$. If $n = 2$, then $\{P \cup Q\}$ is the required set of directed circuits.

Let $n > 2$ and suppose the lemma holds whenever $|VP \cap VQ| < n$. Let b be the last vertex of $VQ - \{x\}$ that is in VP, and let a be the last vertex of VQ incident with an edge of $Q(y, b) - P$. (See Figure 6.) By the inductive hypothesis there exist circuits $C_1, C_2, \ldots, C_{k-1}$ such that

$$\sum_{i=1}^{k-1} C_i = P(a, y) + Q(y, a).$$

The required set of circuits is

$$\{C_1, C_2, \ldots, C_{k-1}, P(x, b) \cup Q(b, x)\},$$



FIGURE 6. P and Q in Lemma 7.

since

$$\begin{aligned} & P(a,y)+Q(y,a)+(P(x,b)\cup Q(b,x)) \\ = \;& P(a,y)+Q(y,a)+P(x,b)+Q(b,x) \\ = \;& P(a,y)+Q(y,a)+P(x,a)+P(a,b)+Q(b,x) \\ = \;& P(x,a)+P(a,y)+Q(y,a)+Q(a,b)+Q(b,x) \\ = \;& P+Q \end{aligned}$$

as $P(a,b)=Q(a,b)$. □

Lemma 8. *Let $\mathcal{C}$ be a set of circuits of even length and empty sum in a directed graph G. Then the parity of the number of clockwise even members of $\mathcal{C}$ is independent of the orientation of G.*

Proof: A change of orientation can be effected by changing orientations of edges one at a time. Each such change leaves unaltered the parity of the number of clockwise even circuits in $\mathcal{C}$. □

Theorem 5. *Let G be a minimal non-Pfaffian near bipartite graph. Let e_1 and e_2 be edges such that $G-\{e_1,e_2\}$ is bipartite and 1-extendible. Let $H=G-\{e_1,e_2\}$, and let f be a 1-factor of H. Let A and B be f-alternating circuits in G of opposite clockwise parity. Suppose that there are exactly two AB-arcs, one containing e_1 and the other e_2. Let the former arc join vertices x_1 and x_2 and the latter vertices y_1 and y_2. Let $A^*=A-\{e_1\}$, and suppose the vertices x_1, y_1, y_2, x_2 appear in that order when A^* is traced from x_1 to x_2. Let H be given its reference orientation with respect to f such that $A^*[x_1,y_1]$ is a directed path from x_1 to y_1. Let S' be a directed path from y_i to x_j and T' a directed path from y_{3-i} to x_{3-j}, where $i,j\in\{1,2\}$. Then G is induced by $A\cup B\cup S'\cup T'$.*

Proof: Without loss of generality we take $i=j=1$. We first show, by using Lemmas 6 and 7, that we can write $A+B$ as a sum of directed circuits included in $A\cup B\cup S'\cup T'$. In order to verify this claim, first we apply Lemma 7 to the directed paths $A(x_1,y_1)$ and S'. Let

$$A(x_1,y_1)+S'=\sum_{V\in\mathcal{V}}V, \tag{1}$$

where $\mathcal{V}$ is a set of directed circuits included in $A(x_1,y_1)\cup S'$. Similarly let

$$A(x_2,y_2)+T'=\sum_{W\in\mathcal{W}}W, \tag{2}$$

where $\mathcal{W}$ is a set of directed circuits included in $A(x_2, y_2) \cup T'$. Now consider the sequence L of edges formed by the edges in the directed path S' followed by those in the directed path $B(x_1, y_2)$. We apply Lemma 6 to L to write

$$S' + B(x_1, y_2) = P + \sum_{X \in \mathcal{X}} X, \tag{3}$$

where P is a directed path from y_1 to y_2 included in $S' \cup B(x_1, y_2)$ and $\mathcal{X}$ is a set of directed circuits included in $S' \cup B(x_1, y_2)$. Similarly we have

$$T' + B(x_2, y_1) = Q + \sum_{Y \in \mathcal{Y}} Y, \tag{4}$$

where Q is a directed path from y_2 to y_1 included in $T' \cup B(x_2, y_1)$ and $\mathcal{Y}$ is a set of directed circuits included in $T' \cup B(x_2, y_1)$. Now we apply Lemma 7 to P and Q to obtain

$$P + Q = \sum_{Z \in \mathcal{Z}} Z, \tag{5}$$

where $\mathcal{Z}$ is a set of directed circuits included in $P \cup Q$. Let $\mathcal{C} = \mathcal{V} + \mathcal{W} + \mathcal{X} + \mathcal{Y} + \mathcal{Z}$. Adding equations (1)–(5) we obtain

$$\begin{aligned} & A(x_1, y_1) + A(x_2, y_2) + B(x_1, y_2) + B(x_2, y_1) \\ = & \sum_{V \in \mathcal{V}} V + \sum_{W \in \mathcal{W}} W + \sum_{X \in \mathcal{X}} X + \sum_{Y \in \mathcal{Y}} Y + \sum_{Z \in \mathcal{Z}} Z \\ = & \sum_{C \in \mathcal{C}} C. \end{aligned}$$

Since the left hand side is $A + B$, $\mathcal{C}$ is the required set of circuits.

In our extended Pfaffian orientation of G, A is the only clockwise even circuit in $\mathcal{C} \cup \{A, B\}$. Therefore by Lemma 8 an odd number of circuits in $\mathcal{C} \cup \{A, B\}$ are clockwise even for any orientation of G. But if there were a Pfaffian orientation of $G[A \cup B \cup \bigcup_{C \in \mathcal{C}} C]$ then every circuit in $\mathcal{C} \cup \{A, B\}$ would be clockwise odd because they are f-alternating. Therefore the graph $G[A \cup B \cup \bigcup_{C \in \mathcal{C}} C]$ is non-Pfaffian, and so $G[A \cup B \cup S' \cup T']$ is non-Pfaffian. By the minimality of G, we deduce that

$$G[A \cup B \cup S' \cup T'] = G.$$

$\square$

Applying this theorem to S and T we find that $G = G[A \cup B \cup S \cup T]$. In fact we chose S and T to satisfy the following definition.

Definition 1. *Let $i, j \in \{1, 2\}$ and let P be a directed path from y_i to x_j under our reference orientation. We say that P is minimal if for every edge $e \in P - (A \cup B)$ there is no directed path from y_i to x_j included in $(A \cup B \cup P) - \{e\}$.*

It is clear that S and T may be assumed to be minimal.

Let P be a directed path from vertex x to vertex y. Let P_1 and P_2 be disjoint subpaths of P such that each edge of P_1 is closer to x in P than is any edge of P_2. In this case we write $P_1 <_P P_2$. If $P_1 = \{a_1\}$ and $P_2 = \{a_2\}$, then we write $a_1 <_P a_2$ instead of $\{a_1\} <_P \{a_2\}$. A similar notation is used for vertices in P.



FIGURE 7. A directed minimal path P from y_i to x_j.

The next lemma is obvious.

Lemma 9. *Let u and v be vertices in $G[A \cup B]$. Under our reference orientation for H there exists at most one directed path in $G[A \cup B]$ from u to v.*

This lemma is used in the proof of the following lemma.

Lemma 10. *(a) Let Q be a minimal directed path from y_i to x_j and let P be a directed path included in $A \cup B$. Let Q_1 and Q_2 be distinct QP-arcs. Then $Q_1 <_Q Q_2$ if and only if $Q_2 <_P Q_1$. (See Figure 7.)*
(b) Conversely let Q' be a directed path from y_i to x_j. If for every directed path P' in $A \cup B$ and every pair of distinct $Q'P'$-arcs Q'_1 and Q'_2 we have $Q'_1 <_{Q'} Q'_2$ if and only if $Q'_2 <_{P'} Q'_1$, then Q' is minimal.

Proof: (a) It suffices to show that if $Q_1 <_Q Q_2$ then $Q_2 <_P Q_1$. Assume the contrary, that is $Q_1 <_Q Q_2$ and $Q_1 <_P Q_2$. Let a be the terminus of Q_1 and b the origin of Q_2. By assumption $P(a, b)$ and $Q(a, b)$ exist, but it are not equal. By Lemma 9 there is an edge $e \in Q(a, b) - (A \cup B)$. The set $Q(y_i, a) \cup P(a, b) \cup Q(b, x_j)$ includes a directed path from y_i to x_j. This path is included in $(A \cup B \cup Q) - \{e\}$, in contradiction to the minimality of Q.

(b) Conversely, assume that Q' is not minimal. Choose $e \in Q' - (A \cup B)$ so that there exists a directed path Q from y_i to x_j in $(A \cup B \cup Q') - \{e\}$. Let u be the last vertex in Q that is also in Q' and satisfies $Q(y_i, u) = Q'(y_i, u)$. Let v be the first vertex in $VQ(u, x_j) - \{u\}$ that is also in Q'. Then $u <_Q v$, $u <_{Q'} v$ and $Q(u, v) \cap Q' = \emptyset$. Hence $Q(u, v) \subseteq A \cup B$ and so $Q(u, v)$ is included in a maximal directed path P' included in $A \cup B$ such that $u <_{P'} v$. Let Q'_1 be the $Q'P'$-arc that includes u and Q'_2 the $Q'P'$-arc that includes v. Then Q'_1 and Q'_2 are distinct, $Q'_1 <_{Q'} Q'_2$ but $Q'_1 <_{P'} Q'_2$. □

3. FORBIDDEN $\overline{A \cup B}$-ARCS

In this section we rule out certain directed $\overline{A \cup B}$-arcs. For that purpose we need the following technical lemma.

Lemma 11. *Let P be a directed $\overline{A \cup B}$-arc. Then there exist $i, j \in \{1, 2\}$ such that P is included in a minimal path Q directed from y_i to x_j.*

Proof: If $S \cap T = \emptyset$, then S and T are vertex disjoint and, since $P \subseteq (S \cup T) - (A \cup B)$ by Theorem 5, it follows that $P \subseteq S$ or $P \subseteq T$. Assume therefore that $S \cap T \neq \emptyset$. Let a and b be, respectively, the first and last vertices of T that are also in S. It follows from Theorem 5 that

$$G = G[A \cup B \cup S \cup T(y_2, a) \cup T(b, x_2)], \tag{6}$$

as there exists a directed path from y_2 to x_2 included in

$$T(y_2, a) \cup S(a, x_1) \cup A(x_1, y_1) \cup S(y_1, b) \cup T(b, x_2).$$

We observe from (6) that any vertex of degree 3 and not in $VA \cup VB$ must be either a or b. It follows that if either a or b were not an internal vertex of P, then P would be included in S or T, since the edges of f incident on a or b are in $S \cap T$. In this case we could choose Q to be S or T. We therefore assume that a and b are the internal vertices of P, since G has no vertices of degree 2.

Let u and v be, respectively, the origin and terminus of P. If $b <_P a$, then $P(u, b) \cup P(a, v) \subseteq S \cap T$, and so $P \subseteq S$ or $P \subseteq T$ according to whether $P(b, a) \subseteq S$ or $P(b, a) \subseteq T$. Therefore we can assume that $a <_P b$. Then $S(a, b) = T(a, b)$. Moreover $P(u, a)$ is included in S or T, and similarly for $P(b, v)$. Without loss of generality we assume that $P(u, a) \subseteq S$. If $P(b, v) \subseteq S$, then we take $Q = S$. Suppose therefore that $P(b, v) \subseteq T$. In this case, we take $Q = S(y_1, b) \cup T(b, x_2)$.

It remains to show that Q is minimal. Suppose not. Then there exists $e \in Q - (A \cup B)$ such that there is a directed path Q' from y_1 to x_2 included in $(A \cup B \cup Q) - \{e\}$. Define $R = T(y_2, b) \cup S(b, x_1)$. Then by Theorem 5 we have $G = G[A \cup B \cup R \cup Q']$. If $e \in S(y_1, a) \cup T(b, x_2)$ then we have the contradiction that $e \notin A \cup B \cup R \cup Q'$. Therefore we suppose that $e \in Q(a, b)$. Then $P \cap Q' = \emptyset$, and we have the contradiction that

$$(S(u, a) \cup T(b, v)) \cap (A \cup B \cup R \cup Q') = \emptyset.$$

□

The next lemma appeared in [4].

Lemma 12. *Let A_1, A_2 be f-alternating circuits in a directed graph G with 1-factor f. Then A_1 and A_2 are of opposite clockwise parity if and only if $A_1 + A_2$ includes an odd number of clockwise even alternating circuits.*

Corollary 1. *The sum $A + B$ is a clockwise even circuit under our extended Pfaffian orientation.*

Proof: Note that

$$A + B = A(x_1, y_1) \cup B(x_2, y_1) \cup A(x_2, y_2) \cup B(x_1, y_2),$$

which is a circuit. Since A and B are of opposite clockwise parity, the result follows from Lemma 12. □

Lemma 13. *Let P be a directed path included in $A \cup B$ such that no internal vertex of P is in $\{x_1, x_2, y_1, y_2\}$. Then there does not exist a directed $\overline{A \cup B}$-arc joining vertices in VP.*

Proof: Suppose there exists a directed $\overline{A \cup B}$-arc Q from $x \in VP$ to $y \in VP$. Then, by Lemma 11, for some $i, j \in \{1, 2\}$ there exists a directed minimal path Z from y_i to x_j that includes Q. By Lemma 5 we may choose a directed path W from y_{3-i} to x_{3-j}; thus $G = G[A \cup B \cup Z \cup W]$ by Theorem 5. There exist a ZP-arc P_1 with terminus x and a ZP-arc $P_2 \neq P_1$ with origin y. Let z_1 be the origin of P_1 and z_2 the terminus of P_2. (See Figure 8.) Since $x <_Z y$ we have $P_1 <_Z P_2$, so that $P_2 <_P P_1$ by Lemma 10. Therefore $z_2 <_P z_1$.

Let C be the circuit $Q \cup P(y, x)$. First we show that we may assume there to be at most one CW-arc. Suppose there are two such arcs, W_1 and W_2, where $W_1 <_W W_2$. Let a be the terminus of W_1 and b the origin of W_2. Let W^* be a directed path from y_{3-i} to x_{3-j} included in

$$W(y_{3-i}, a) \cup C(a, b) \cup W(b, x_{3-j}).$$

The number of CW^*-arcs is less than the number of CW-arcs. By repeating the argument if necessary and appealing to the finiteness of G, we may therefore assume that there is at most one CW-arc. If such an arc exists, let its origin be w_1 and its terminus w_2. (See Figure 8.) We also note that there is a unique CZ-arc, by the minimality of Z.

Let f' be the 1-factor $f + C$, and let $A' = A + C$, $B' = B + C$, $Z' = Z + C$ and $W' = W + C$. Now we show that

$$G = G[A' \cup B' \cup Z' \cup W']. \tag{7}$$

By Lemma 12, A' and B' are f'-alternating circuits containing e_1 and e_2 of opposite clockwise parity, since A and B are of opposite clockwise parity and $A' + B' = A + B$. Moreover there are exactly two $A'B'$-arcs and the vertices of degree 3 in $G[A' \cup B']$ are the same as those in $G[A \cup B]$, since

$$\{x_1, x_2, y_1, y_2\} \cap (VC - \{x, y\}) = \emptyset.$$

In addition Z' would become a directed path from y_i to x_j if C were reoriented, and a similar statement holds for W'. Thus (7) holds, by Theorem 5 with f, A and B replaced by f', A' and B' respectively.

Now we observe that

$$(A' \cup B') \cap P(y, x) = \emptyset$$

since

$$\{x_1, x_2, y_1, y_2\} \cap (VP - \{x, y\}) = \emptyset.$$

Note that

$$Z' = Z(y_i, z_1) \cup P(z_2, z_1) \cup Z(z_2, x_j).$$

Therefore

$$Z' \cap (P_1 \cup P_2) = \emptyset.$$

Thus $(A' \cup B' \cup Z') \cap (P_1 \cup P_2) = \emptyset$, and so $P_1 \cup P_2 \subseteq W'$ by (7). We deduce that w_1 and w_2 exist.

Next we show that either $w_1, w_2 \in VP(z_2, z_1)$ and $z_2 <_P w_1 <_P w_2 <_P z_1$, or $w_1, w_2 \in VQ$ and $w_1 <_Q w_2$. First,

$$W' = W(y_{3-i}, w_1) \cup C(w_2, w_1) \cup W(w_2, x_{3-j}).$$

Hence

$$P_1 \cup P_2 \subseteq C(w_2, w_1),$$

and the desired conclusion follows.

Case 1: Suppose first that $w_1, w_2 \in VP(z_2, z_1)$ and $z_2 <_P w_1 <_P w_2 <_P z_1$. After reorientation of C, let X be a directed path from y_i to x_{3-j} included in

$$Z(y_i, z_1) \cup C(z_1, w_2) \cup W(w_2, x_{3-j})$$

and let Y be a directed path from y_{3-i} to x_j included in

$$W(y_{3-i}, w_1) \cup C(w_1, z_2) \cup Z(z_2, x_j).$$

Thus $G = G[A' \cup B' \cup X \cup Y]$. We now have the contradiction that

$$(P_1 \cup C(w_2, w_1) \cup P_2) \cap (A' \cup B' \cup X \cup Y) = \emptyset.$$

Case 2: Suppose on the other hand that $w_1, w_2 \in VQ$ and $w_1 <_Q w_2$. Without loss of generality we may assume that P is a maximal directed path in $A \cup B$ such that no internal vertex is in $\{x_1, x_2, y_1, y_2\}$. Let P be directed from vertex u to vertex v. Thus $u \in \{u_1, u_2, x_1, x_2, y_1, y_2\}$ and $v \in \{x_1, x_2, y_1, y_2, v_1, v_2\}$.

First we show that $u \neq y_i$. If $u = y_i$ then we observe that there is a directed path Z^* from y_i to x_j included in $P(u, z_2) \cup Z(z_2, x_j)$. Therefore

$$(A \cup B \cup Z^* \cup W) \cap (C(x, w_1) \cup C(w_2, y)) = \emptyset.$$

By Theorem 5 we have the contradiction that $G = G[A \cup B \cup Z^* \cup W]$. Thus $u \neq y_i$. A similar argument, with Z^* included in $Z(y_i, z_1) \cup P(z_1, v)$, shows that $v \neq x_j$.

Next we show that $u \neq y_{3-i}$. Otherwise we define W^* to be a directed path from y_{3-i} to x_{3-j} included in

$$P(u, x) \cup C(x, w_2) \cup W(w_2, x_{3-j}).$$

The union $A \cup B \cup Z \cup W^*$ does not contain the edge of $W - Q$ incident on w_1. This result contradicts Theorem 5, since $G = G[A \cup B \cup Z \cup W^*]$. Thus $u \neq y_{3-i}$. A similar argument, with W^* included in

$$W(y_{3-i}, w_1) \cup C(w_1, y) \cup P(y, v),$$

shows that $v \neq x_{3-j}$.

Now we show that $u \neq x_{3-j}$. Otherwise we define Z^* as a directed path from y_i to x_j included in

$$Z(y_i, w_2) \cup W(w_2, u) \cup P(u, z_2) \cup Z(z_2, x_j).$$

Then

$$(A \cup B \cup Z^* \cup W) \cap C(w_2, y) = \emptyset,$$

in contradiction to Theorem 5 since $G = G[A \cup B \cup Z^* \cup W]$. A similar argument shows that $v \neq y_{3-i}$.

Since $v \notin \{x_1, x_2\}$ we have $u \notin \{u_1, u_2\}$. We conclude that $u = x_j$, and similarly $v = y_i$. Define $P' = P + C$. Remember that W' is the only member of $\{A', B', W', Z'\}$ meeting $P_1 \cup P_2$. If C is reoriented then there is a directed path W'' from y_{3-i} to x_{3-j} included in

$$W'(y_{3-i}, w_1) \cup P'(w_1, v) \cup Z'(v, u) \cup P'(u, w_2) \cup W'(w_2, x_{3-j}).$$



FIGURE 8. The situation in Lemma 13.

Thus

$$W'' \cap (P_1 \cup P_2) = \emptyset.$$

We now have a contradiction, since $G = G[A' \cup B' \cup Z' \cup W'']$. ☐

Lemma 14. *There is no directed* $\overline{A \cup B}$*-arc joining vertices in distinct* $A\overline{B}$*-arcs, or in distinct* $B\overline{A}$*-arcs.*

Proof: In view of the symmetry between A and B it suffices to prove that no $\overline{A \cup B}$-arc is directed from a vertex in $A(x_1, y_1)$ to a vertex in $A(x_2, y_2)$. Suppose such an arc P exists, joining a vertex $x'_1 \in VA(x_1, y_1)$ to a vertex $y'_2 \in VA(x_2, y_2)$. (See Figure 9.) Let

$$B' = A(u_1, x'_1) \cup P \cup A(y'_2, v_2) \cup \{e_2\} \cup B(u_2, v_1) \cup \{e_1\}.$$

This is an f-alternating circuit containing e_1 and e_2. Observe that $B(x_1, y_2)$ is the only $B\overline{B'}$-arc. It follows that B and B' have the same clockwise parity, for otherwise A and B could have been chosen to have a unique AB-arc. Henceforth B' will play the rôle previously assumed by B. The circuit A will play the same rôle as before, but we define $x'_2 = x_2$ and $y'_1 = y_1$. Note that there are exactly two AB'-arcs, one containing e_1 and the other containing e_2 and that $B(x_1, y_2)$ is an $\overline{A \cup B'}$-arc. Therefore, by Lemma 11, for some $i, j \in \{1, 2\}$ there exists a directed minimal path X from y'_i to x'_j including $B(x_1, y_2)$.

We now show that $i = 1$ and $j = 2$. Included in the set $X(y'_i, x_1) \cup A(x_1, x'_1)$ is a directed path W from y'_i to x'_1. This path is included in $(A \cup B \cup X) - B(x_1, y_2)$, in contradiction to the minimality of X if $j = 1$. Therefore $j = 2$. Similarly, included in the set $A(y'_2, y_2) \cup X(y_2, x'_2)$ is a directed path Z from y'_2 to x'_2. This path is included in $(A \cup B \cup X) - B(x_1, y_2)$, in contradiction to the minimality of X if $i = 2$. Therefore $i = 1$.

By Theorem 5 we have $G = G[A \cup B' \cup W \cup Z]$, in contradiction to the fact that

$$B(x_1, y_2) \cap (A \cup B' \cup W \cup Z) = \emptyset.$$

☐



FIGURE 9. P is a directed arc joining a vertex in $A(x_1, y_1)$ to a vertex in $A(x_2, y_2)$.

Lemma 15. *There is no directed $\overline{A \cup B}$-arc joining a vertex in an $A\overline{B}$-arc to a vertex in a $B\overline{A}$-arc.*

Proof: By symmetry it suffices to prove that no $\overline{A \cup B}$-arc is directed from a vertex in $A(x_1, y_1)$ to a vertex in $B(x_2, y_1)$. Suppose such an arc P exists, joining a vertex $v \in VA(x_1, y_1)$ to a vertex $y_1' \in VB(x_2, y_1)$. (See Figure 10.) Let

$$A' = A(u_1, v) \cup P \cup B(y_1', v_1) \cup \{e_2\} \cup A(u_2, v_2) \cup \{e_1\}.$$

This is an f-alternating circuit containing e_1 and e_2. Observe that $A(v, y_1)$ is the only $A\overline{A'}$-arc. It follows that A and A' have the same clockwise parity. Henceforth A' will play the rôle previously assumed by A. (See Figure 10, second picture, where A' is drawn as a circle.) The circuit B will play the same rôle as before, but we define $x_1' = x_1$, $x_2' = x_2$ and $y_2' = y_2$. Note that $A(v, y_1)$ is an $\overline{A' \cup B}$-arc. Therefore, by Lemma 11, for some $i, j \in \{1, 2\}$ there exists a directed minimal path X from y_i' to x_j' including $A(v, y_1)$. Let P_1 be the $A'X$-arc with terminus v and P_2 be the $A'X$-arc with origin y_1. Then $P_1 <_X P_2$ and $P_1 <_{A'(u_1, v_1)} P_2$ in contradiction to the minimality of X. $\square$

Lemma 16. *For each $i, j \in \{1, 2\}$ there is no directed $\overline{A \cup B}$-arc from a vertex in $A(y_i, v_i)$ to a vertex in $A(u_j, x_j)$.*

Proof: It suffices to prove the lemma for $i = j = 1$. Suppose such a directed arc P exists. Let P be directed from vertex v to vertex u. Define

$$C = A(u, v) \cup P.$$

Let $f' = f + C$,

$$\begin{aligned} A' &= A + C \\ &= P \cup A(v, v_1) \cup \{e_2\} \cup A(u_2, v_2) \cup \{e_1\} \cup A(u_1, u) \end{aligned}$$



FIGURE 10. P is a directed arc joining a vertex in $A(x_1, y_1)$ to a vertex in $B(x_2, y_1)$.

and

$$\begin{aligned} B' &= B + C \\ &= P \cup B(v, v_1) \cup \{e_2\} \cup B(x_1, v_2) \cup A(x_1, y_1) \cup B(u_2, y_1) \cup \{e_1\} \cup B(u_1, u). \end{aligned}$$

Thus A' and B' are f'-alternating circuits of opposite clockwise parity containing e_1 and e_2. However $A(x_2, y_2)$ is the only $A'\overline{B'}$-arc. This result contradicts the assumption that for every choice of A and B there are at least two AB-arcs. $\square$

Lemma 17. *Let P be a directed $A\overline{B}$-arc or a directed $B\overline{A}$-arc and let Q be a directed AB-arc in H having neither end in common with an end of P. Then there is no pair X, Y of $\overline{A \cup B}$-arcs such that X is directed from a vertex $u \in VP$ to a vertex $v \in VQ$, Y is directed from a vertex $w \in VQ$ to a vertex $x \in VP$, $x <_P u$ and $v <_Q w$. (See Figure 11.)*

Proof: By the symmetry between A and B we may assume that $P = A(x_1, y_1)$. Therefore $Q = A(u_2, x_2)$ or $Q = A(y_2, v_2)$. By symmetry we may assume the latter case obtains.

Suppose X and Y exist. Let

$$C = A(v, w) \cup Y \cup A(x, u) \cup X.$$

Define $f' = f + C$,

$$\begin{aligned} A' &= A + C \\ &= A(w, v_2) \cup \{e_2\} \cup A(u, v_1) \cup X \cup A(u_2, v) \cup \{e_1\} \cup A(u_1, x) \cup Y \end{aligned}$$

and

$$\begin{aligned} B' &= B + C \\ &= B(w, v_2) \cup \{e_2\} \cup B(u_2, v_1) \cup \{e_1\} \cup B(u_1, v) \cup X \cup A(x, u) \cup Y. \end{aligned}$$



FIGURE 11. The situation in Lemma 17 before and after reorienting C.

Then A' and B' are f'-alternating circuits containing e_1 and e_2 and having opposite clockwise parity. Reorient C and define

$$D = B'(u_2, v_1) \cup \{e_2\} \cup A'(u_1, v_2) \cup \{e_1\}.$$

(See Figure 11.) Then D is f'-alternating and contains e_1 and e_2. If D is of opposite clockwise parity to A' then we have a contradiction because there is a unique $A'\overline{D}$-arc $A'(x_2, y_1)$; otherwise B' and D have opposite clockwise parity and there is another contradiction since $B'(x_1, x)$ is the only $B'\overline{D}$-arc. $\square$

4. PRODUCTION OF A LIST OF CASES TO CONSIDER

We now introduce a notation to describe a minimal directed path X from y_i to x_j for $i, j \in \{1, 2\}$. Traversed from y_i to x_j, X meets a succession of directed $(A \cup B)X$-arcs in H. The *trace* of X is the sequence obtained from X by recording: 0 for each $A(x_j, y_i)X$-arc, $0'$ for each $A(x_{3-j}, y_{3-i})X$-arc, 1 for each $B(x_{3-j}, y_i)X$-arc, $1'$ for each $B(x_j, y_{3-i})X$-arc, 2 for each $(A \cap B)(y_{3-i}, v_{3-i})X$-arc, $2'$ for each $(A \cap B)(u_{3-j}, x_{3-j})X$-arc.

Figure 12 shows an example of a directed minimal path from y_1 to x_1 with trace $02110'1'02'1$. By Lemma 10(a) the graph $G[A \cup B \cup X]$ is determined up to homeomorphism by the trace of X. In particular, there are a unique $A(y_i, v_i)X$-arc and a unique $A(u_j, x_j)X$-arc.

Lemma 18. *Let W be a string over $\{0, 0', 1, 1', 2, 2'\}$.*
(a) It is possible to choose A, B, f, x_1, x_2, y_1, y_2 and a directed minimal path X from y_1 to x_1 in G such that the trace of X is $0W$ if and only if it is possible, without altering $A \cup B \cup X$, to choose A, B, f, x_1, x_2, y_1, y_2 and a directed minimal path X from y_1 to x_1 in G such that the trace of X is $1W$.
(b) It is possible to choose A, B, f, x_1, x_2, y_1, y_2 and a directed minimal path X from y_1 to x_1 in G such that the trace of X is $W0$ if and only if it is possible, without altering



FIGURE 12. A directed minimal path with trace $02110'1'02'1$.

$A \cup B \cup X$, to choose A, B, f, x_1, x_2, y_1, y_2 and a directed minimal path X from y_1 to x_1 in G such that the trace of X is $W1'$.

Proof: By symmetry, it suffices to prove (a). Suppose the trace of X is $0W$. There is an $A(x_1, y_1)X$-arc that corresponds to the first 0 in the trace of X. Let v be its origin, and let

$$C = X(y_1, v) \cup A(v, y_1).$$

Let u be the terminus of the $A(y_1, v_1)X$-arc and w the terminus of the $A(x_1, y_1)X$-arc with origin v. (See Figure 13.) Define $f' = f + C$,

$$\begin{aligned} A' &= A + C \\ &= C(u, v) \cup A(u_1, v) \cup \{e_1\} \cup A(u_2, v_2) \cup \{e_2\} \cup A(u, v_1), \end{aligned}$$

$$\begin{aligned} B' &= B + C \\ &= C(u, y_1) \cup B(u_2, y_1) \cup \{e_1\} \cup B(u_1, v_2) \cup \{e_2\} \cup B(u, v_1) \end{aligned}$$

and

$$\begin{aligned} X' &= X + A(v, y_1) \\ &= C(w, v) \cup X(w, x_1). \end{aligned}$$

Then A' and B' are f'-alternating circuits containing e_1 and e_2 and having opposite clockwise parity. Moreover there are exactly two $A'B'$-arcs, one containing e_1 and the other containing e_2, and the vertices of degree 3 in $G[A' \cup B']$ are x_1, x_2, v, y_2. After reorientation of C, X' is a directed path from v to x_1. The trace of X' is



FIGURE 13. The situation in Lemma 18 before and after reorienting C.

$1W$, as $B'(y_1, w)$ is a $B'(x_2, v)X'$-arc which replaces the $A(x_1, y_1)X$-arc $A(v, w)$. (See Figure 13.) Moreover X' is minimal: X' satisfies the condition in Lemma 10(b) since X does and

$$X'(v, w) \cap (A' \cup B') = C(v, u) \cup C(y_1, w).$$

For the converse in (a) note that $f = f' + C$, $A = A' + C$, $B = B' + C$ and $X = X' + A(v, y_1)$. ☐

Lemma 19. *For some choice of A, B, x_1, x_2, y_1, y_2, f there is a directed minimal path S' from y_1 to x_1 with trace 0 or $0'$. (See Figure 14.)*

Proof: We choose A, B, x_1, x_2, y_1, y_2, f and a directed path S' from y_1 to x_1 so that $A \cup B \cup S'$ is minimal. Thus S' is minimal.

Suppose the trace of S' contains 2. There is an $(A \cap B)(y_2, v_2)S'$-arc; let v be its terminus. Included in the set $A(y_2, v) \cup S'(v, x_1)$ is a directed path from y_2 to x_1, in contradiction to the minimality of $A \cup B \cup S'$ since $S'(y_1, v) - (A \cup B) \neq \emptyset$. Therefore the trace of S' does not contain 2, and similarly does not contain $2'$.

The trace of S' contains none of 00, 11, $0'0'$, $1'1'$ by Lemma 13, none of $00'$, $0'0$, $11'$, $1'1$ by Lemma 14, none of 01, $01'$, $0'1$, $0'1'$, 10, $10'$, $1'0$, $1'0'$ by Lemma 15, and is non-empty by Lemma 16. We infer that the trace of S is one of 0, $0'$, 1, $1'$. By Lemma 18 the case that the trace of S' is 1 or $1'$ can be reduced to the case that the trace of S' is 0. ☐

Because of this lemma we henceforth assume that the trace of S is 0 or $0'$. Given this choice for the trace of S, we now turn our attention to the trace of T.

In the following we produce a finite list of possible traces of T and therefore a finite list of graphs we will consider in the following section. In order to do so we distinguish the two cases $S \cap T = \emptyset$ and $S \cap T \neq \emptyset$. First we assume that $S \cap T = \emptyset$.

Lemma 20. *Let W be a string over $\{0, 0', 1, 1', 2, 2'\}$, and let $S \cap T = \emptyset$.*
(a) Suppose that the trace of T is $1W$. Then there exist A, B, f, x_1, x_2, y_1, y_2, a



FIGURE 14. A directed minimal path S' with trace 0 and with trace $0'$.

directed minimal path from S' y_1 *to* x_1 *with trace* 0 *or* $0'$ *and a directed minimal path* T' *from* y_2 *to* x_2 *with trace* $0W$.
(b) Suppose that the trace of T *is* $W1'$. *Then there exist* A, B, f, x_1, x_2, y_1, y_2, *a directed minimal path* S' *from* y_1 *to* x_1 *with trace* 0 *or* $0'$ *and a directed minimal path* T' *from* y_2 *to* x_2 *with trace* $W0$.

Proof: That T' exists follows by a proof similar to that of the corresponding assertion in Lemma 18. The reorientation of the corresponding f-alternating circuit C does not affect S, since $S \cap C = \emptyset$. Therefore we may take $S' = S$. □

Thus we can assume that the first symbol in the trace of T is not 1 and that the last symbol in the trace of T is not $1'$. In the following we will refer to this property of T as (A).

Lemma 21. *Suppose* $S \cap T = \emptyset$ *and the trace of* T *contains one of* $20'$, $21'$, $12'$, $0'2'$. *Then there exist* A', B', f', *a directed minimal path* S' *from* y_1 *to* x_1 *with trace* 0 *or* $0'$ *and a directed minimal path* T' *from* y_2 *to* x_2 *such that* $S' \cap T' \neq \emptyset$.

Proof: It suffices to consider the case where the trace of T contains $20'$, for the other cases are similar. In this case there is a $T\overline{A \cup B}$-arc with origin $u \in VA(y_1, v_1)$ and terminus $v \in VA(x_1, y_1)$. Let u' be the origin of the $A(y_1, v_1)T$-arc with terminus u, and v' the terminus of the $A(x_1, y_1)T$-arc with origin v. Let w be the terminus of the unique $A(y_1, v_1)S$-arc. (See Figure 15.)

Since $S \cap T = \emptyset$ we have $w <_{A(y_1,v_1)} u'$. If S has trace 0, then let x be the terminus of the unique $A(x_1, y_1)S$-arc and x' its origin. We have $x <_{A(x_1,y_1)} v$: otherwise if we define

$$S^* = A(y_1, u) \cup T(u, v) \cup A(v, x) \cup S(x, x_1)$$

then $G = G[A \cup B \cup S^* \cup T]$ by Theorem 5, in contradiction to the fact that

$$S(w, x') \cap (A \cup B \cup S^* \cup T) = \emptyset.$$



Figure 15. The situation in Lemma 21 before and after reorienting C.

In any case, define

$$C = T(u, v) \cup A(v, u).$$

This is an f-alternating circuit such that $C \cap S = A(y_1, w)$. Reorient C and define $f' = f + C$,

$$\begin{aligned} A' &= A + C \\ &= C(v, u) \cup A(u_1, v) \cup \{e_1\} \cup A(u_2, v_2) \cup \{e_2\} \cup A(u, v_1), \end{aligned}$$

$$\begin{aligned} B' &= B + C \\ &= C(y_1, u) \cup B(u_2, y_1) \cup \{e_1\} \cup B(u_1, v_2) \cup \{e_2\} \cup B(u, v_1), \end{aligned}$$

$$\begin{aligned} S' &= S + C(v, y_1) \\ &= C(v, w) \cup S(w, x_1) \end{aligned}$$

and

$$\begin{aligned} T' &= T + C \\ &= T(y_2, u') \cup C(u', v') \cup T(v', x_2). \end{aligned}$$

Then A' and B' are f'-alternating circuits containing e_1 and e_2 and having opposite clockwise parity. There are exactly two $A'B'$-arcs, the vertices of degree 3 in $G[A' \cup B']$ are x_1, x_2, v and y_2, S' is a directed path from v to x_1 and T is a directed path from y_2 to x_2 (see Figure 15). Moreover S and S' have equal trace and $S' \cap T' = C(u', w) \neq \emptyset$. Finally S' and T' are minimal: both satisfy the condition in Lemma 10(b) since $S'(v, w) \cap (A' \cup B') = C(v, u)$ and $T'(u', v') \cap (A' \cup B') = C(y_1, v')$. $\square$

Thus we assume that the trace of T contains none of $20'$, $21'$, $12'$, $0'2'$, if $S \cap T = \emptyset$. In the following we will refer to this property of T as (B).

The following lemma gives a complete list of graphs to be considered if $S \cap T = \emptyset$. We use $*$ to denote an arbitrary string of symbols, and Λ to denote the empty string.

Lemma 22. *Suppose $S \cap T = \emptyset$ and that T has properties (A) and (B). Then the trace of T is one of* 0, $0'$, 20, 21, $02'$, $1'2'$.

Proof: First we see that the symbols in the trace of T alternate between the sets $\{0, 0', 1, 1'\}$ and $\{2, 2'\}$, for the trace of T contains none of 00, 11, 22, $0'0'$, $1'1'$, $2'2'$ by Lemma 13, none of $00'$, $0'0$, $11'$, $1'1$ by Lemma 14, none of 01, $01'$, $0'1$, $0'1'$, 10, $10'$, $1'0$, $1'0'$ by Lemma 15, and does not contain $22'$ by Lemma 16 or $2'2$ by Lemma 10(a).

Next we show that the trace of T does not contain both 2 and $2'$. Suppose that the trace of T contains $2' * 2$. Thus $x_1, y_1 \in VT$ by Lemma 10(a). This is a contradiction to $S \cap T = \emptyset$. Suppose that the trace of T contains $2 * 2'$. Choose 2 and $2'$ in the trace of T, with the chosen $2'$ appearing later than the chosen 2. Let u be the terminus of the $A(y_1, v_1)T$-arc that corresponds to the chosen 2 and let v be the origin of the $A(u_1, x_1)T$-arc that corresponds to the chosen $2'$. There is a directed path from y_1 to x_1 included in

$$A(y_1, u) \cup T(u, v) \cup A(v, x_1).$$

By Theorem 5 we have $G = G[A \cup B \cup T]$, in contradiction to

$$(S - (A \cup B)) \cap (A \cup B \cup T) = \emptyset.$$

We distinguish the following cases:

1. The trace of T contains neither 2 nor $2'$.
2. The trace of T contains 2 and consequently does not contain $2'$.
3. The trace of T contains $2'$ and consequently does not contain 2.

Case 1: By Lemma 16 the trace is not empty and therefore the trace of T is one of 0, $0'$, 1, $1'$ in this case. By (A) the case that the trace is 1 or $1'$ is not possible.

Case 2: The symbols in the trace of T alternate between the sets $\{2\}$ and $\{0, 0', 1, 1'\}$. By (B) and Lemma 16 every 2 in the trace must be immediately followed by 0 or 1. Therefore the trace contains at most one 2 by Lemma 17 and consequently exactly one 2. In fact the trace of T is $x2y$, where $x \in \{\Lambda, 0, 0', 1, 1'\}$ and $y \in \{0, 1\}$.

We show that $x = \Lambda$. We have $x \notin \{0', 1'\}$, for otherwise $y_1 \in VS \cap VT$ by Lemma 10(a). Suppose $x = 0$. If $y = 0$, we have a contradiction by Lemma 17. If $y = 1$, we have a contradiction by Lemma 17 also, since in this case there exist A', B', f', x_1', x_2', y_1', y_2' in G and a directed minimal path T' from y_2' to x_2' with trace 121 by Lemma 18. Similarly we obtain a contradiction if we suppose that $x = 1$. Therefore the trace of T is either 20 or 21, if the trace contains 2.

Case 3: Similarly the trace of T is either $02'$ or $1'2'$, if it contains $2'$. □

Remark 1. *The case where the trace of T is either* $02'$ *or* $1'2'$ *can be reduced to the case where the trace of T is either* 20 *or* 21. *In order to see this suppose that the trace of T is either* $02'$ *or* $1'2'$ *and switch to the reference orientation with respect to (N, M, f). Then S is a directed path from $y_1' = x_1$ to $x_1' = y_1$ with trace* 0 *or* $0'$ *and T is a directed path from $y_2' = x_2$ to $x_2' = y_2$ with trace* 20 *or* 21.

Now we consider the case where $S \cap T \neq \emptyset$. First we prove the following consequence of Theorem 5.



FIGURE 16. There are two possible cases: either $a <_S b$ or $b <_S a$.

Corollary 2. *Let S' be a directed path from y_1 to x_1, T_1 a directed path from y_2 to a vertex in S' and T_2 a directed path from a vertex in S' to x_2. Then*

$$G = G[A \cup B \cup S' \cup T_1 \cup T_2].$$

Proof: By Theorem 5 we have to show that there exists a directed path from y_2 to x_2 in $G[A \cup B \cup S' \cup T_1 \cup T_2]$. Let a be the terminus of T_1 and b the origin of T_2. Then such a path is included in

$$T_1 \cup S(a, x_1) \cup A(x_1, y_1) \cup S(y_1, b) \cup T_2.$$

$\square$

Since $S \cap T \neq \emptyset$, there exists a first vertex a in T that is also in S, and a last vertex b in T that is also in S. (See Figure 16.) Let $T_1 = T(y_2, a)$ and $T_2 = T(b, x_2)$. By Corollary 2,

$$G = G[A \cup B \cup S \cup T_1 \cup T_2].$$

We define the trace of T_1 and T_2 in a manner analogous to the definition of the trace of a directed minimal path from y_2 to x_2. Note that T_1 and T_2 satisfy the condition in Lemma 10(b), since they are directed subpaths of the directed minimal path T.

Lemma 23. *Let W be a string over $\{0, 0', 1, 1', 2, 2'\}$, and let $S \cap T \neq \emptyset$.*
(a) Suppose that the trace of T_1 is $1W$. Then there exist A, B, f, x_1, x_2, y_1, y_2, a directed minimal path from y_1 to x_1 with trace 0 or $0'$ and a directed minimal path T from y_2 to x_2 such that the trace of T_1 is $0W$.
(b) Suppose that the trace of T_2 is $W1'$. Then there exist A, B, f, x_1, x_2, y_1, y_2, a directed minimal path from y_1 to x_1 with trace 0 or $0'$ and a directed minimal path T from y_2 to x_2 such that the trace of T_2 is $W0$.

Proof: Similar to the proof of Lemma 20. $\square$

Therefore we can assume that the first symbol in the trace of T_1 is not 1 and that the last symbol in the trace of T_2 is not $1'$. In the following we will refer to this property of T_1 and T_2 as (A').

In the directed path S there are exactly 6 vertices of degree 3 in $G[A \cup B \cup S]$, the first being y_1 and the last being x_1. We label the other such vertices w_1, w_2, w_3 and w_4 in the order they occur when we traverse S from y_1 to x_1. (See Figure 14.) Note that $a \notin \{w_1, w_3, x_1\}$ and $b \notin \{y_1, w_2, w_4\}$, since the vertices of G have indegree 1 or outdegree 1.

Lemma 24. *The vertices a and b are not both in $VS(y_1, w_1)$ and not both in $VS(w_4, x_1)$.*

Proof: By symmetry it suffices to show that $\{a, b\} \not\subseteq VS(y_1, w_1)$. Suppose the contrary.

First we assume that the trace of S is 0. We define

$$C = S(w_1, w_2) \cup A(w_2, w_1).$$

This is an f-alternating circuit. Furthermore we define $f' = f + C$,

$$\begin{aligned} A' &= A + C \\ &= C(w_1, w_2) \cup A(u_1, w_2) \cup \{e_1\} \cup A(u_2, v_2) \cup \{e_2\} \cup A(w_1, v_1), \end{aligned}$$

$$\begin{aligned} B' &= B + C \\ &= C(w_1, y_1) \cup B(u_2, y_1) \cup \{e_1\} \cup B(u_1, v_2) \cup \{e_2\} \cup B(w_1, v_1). \end{aligned}$$

By Lemma 12, A' and B' are f'-alternating circuits containing e_1 and e_2, of opposite clockwise parity, such that there are exactly two $A'B'$-arcs. The vertices of degree 3 in $G[A' \cup B']$ are x_1, x_2, w_2, y_2.

First we assume $a <_S b$. In this case $a = y_1$ and $b = w_1$, for otherwise we would have vertices of degree 2. We define the paths

$$X' = T_1 \cup C(w_3, a) \cup S(w_3, x_1)$$

and

$$Y' = C(b, w_2) \cup T_2.$$

These paths would become directed from y_2 to x_1 and from w_2 to x_2, respectively, if C were reoriented. Therefore

$$G = G[A' \cup B' \cup X' \cup Y']$$

by Theorem 5, in contradiction to

$$C(a, b) \cap (A' \cup B' \cup X' \cup Y') = \emptyset.$$

Now we assume that $b <_S a$. We reorient C and define the directed path

$$\begin{aligned} T' &= T + C \\ &= T_1 \cup C(a, b) \cup T_2 \end{aligned}$$

and a directed path S' from w_2 to x_1 included in

$$C(w_2, b) \cup T_2 \cup B(x_2, y_1) \cup C(y_1, w_3) \cup S(w_3, x_1).$$

By Theorem 5

$$G = G[A' \cup B' \cup S' \cup T'],$$

in contradiction to

$$C(b, y_1) \cap (A' \cup B' \cup S' \cup T') = \emptyset.$$

Next we consider the case that the trace of S is $0'$. We define

$$D = S(w_1, w_4) \cup A(w_4, w_1).$$

This is an f-alternating circuit. Furthermore we define $f'' = f + D$,

$$\begin{aligned} A'' &= A + D \\ &= D(w_1, w_2) \cup A(u_2, w_2) \cup \{e_1\} \cup A(u_1, w_4) \cup \\ &\quad D(w_3, w_4) \cup A(w_3, v_2) \cup \{e_2\} \cup A(w_1, v_1), \end{aligned}$$

$$\begin{aligned} B'' &= B + D \\ &= D(w_1, w_4) \cup B(u_1, w_4) \cup \{e_1\} \cup B(u_2, y_1) \cup \\ &\quad D(x_1, y_1) \cup B(x_1, v_2) \cup \{e_2\} \cup B(w_1, v_1). \end{aligned}$$

By Lemma 12, A'' and B'' are f''-alternating circuits including e_1 and e_2, of opposite clockwise parity, such that there are exactly two $A''B''$-arcs. The vertices of degree 3 in $G[A'' \cup B'']$ are w_3, x_2, w_2, y_2.

First we assume $a <_S b$. Again we have $a = y_1$ and $b = w_1$, for otherwise G would have vertices of degree 2. We define the paths

$$X'' = D(b, w_2) \cup T_2$$

and

$$Y'' = T_1 \cup D(w_3, a).$$

These paths would become directed from w_2 to x_2 and from y_2 to w_3, respectively, if D were reoriented. Therefore

$$G = G[A'' \cup B'' \cup X'' \cup Y'']$$

by Theorem 5, in contradiction to

$$D(a, b) \cap (A'' \cup B'' \cup X'' \cup Y'') = \emptyset.$$

Now we assume that $b <_S a$. We reorient D and define the directed path

$$\begin{aligned} T'' &= T + D \\ &= T_1 \cup D(a, b) \cup T_2 \end{aligned}$$

and a directed path S'' from w_2 to w_3 included in

$$D(w_2, b) \cup T_2 \cup B(x_2, y_1) \cup D(y_1, w_3).$$

By Theorem 5

$$G = G[A'' \cup B'' \cup S'' \cup T''],$$

in contradiction to

$$D(b, y_1) \cap (A'' \cup B'' \cup S'' \cup T'') = \emptyset.$$

□

Lemma 25. *The traces of* T_1 *and* T_2 *are either empty or* 0.

Proof: By symmetry it suffices to show that the trace of T_1 is either empty or 0. First we show that the edge in T_1 incident on a is not in $A \cup B$. This edge exists, for $T_1 \neq \emptyset$ since the trace of S is 0 or $0'$. Let a^* be its origin and suppose that $(a^*, a) \in A \cup B$. Note that $(a^*, a) \notin f$ because S is f-alternating. Thus the edge of f incident on a^* is in T_1 and in $A \cup B$. Now we have the contradiction that a^* is a vertex of degree 2, since $a^* \notin VS \cup VT_2$.

We use this observation to show that the trace of T_1 contains none of $0'$, $1'$, $2'$.

Suppose that the trace of T_1 contains $0'$. Let a' be the origin of an $A(x_1, y_1)T_1$-arc. Clearly $a' \neq a$. Included in

$$T_1(y_2, a') \cup A(a', y_1)$$

is a directed path T_1' from y_2 to a vertex in S with

$$T_1' \cap (T_1(a', a) - (A \cup B)) = \emptyset.$$

By Corollary 2 we have

$$G = G[A \cup B \cup S \cup T_1' \cup T_2].$$

Since

$$(T_1(a', a) - (A \cup B)) \cap (A \cup B \cup S \cup T_1' \cup T_2) = \emptyset,$$

it follows that

$$T_1(a', a) - (A \cup B) = \emptyset.$$

Therefore we have the contradiction that the edge of T_1 incident on a is in $A \cup B$. The proof that the trace of T_1 contains neither $1'$ nor $2'$ is similar. Likewise the trace of T_2 contains none of $0'$, 1 and 2.

Next we show that the trace of T_1 does not contain 2. Assume the contrary and let v be the terminus of the last $A(y_1, v_1)T_1$-arc. Then there is a directed path S' from y_1 to x_1 included in

$$A(y_1, v) \cup T_1(v, a) \cup S(a, x_1)$$

with

$$S' \cap (S(y_1, a) - (A \cup B)) = \emptyset.$$

Therefore

$$S(y_1, a) \subseteq A \cup B \cup T,$$

since $G = G[A \cup B \cup S' \cup T]$ by Theorem 5. Consequently,

$$a \notin (VS(w_1, w_2) - \{w_1\}) \cup (VS(w_3, w_4) - \{w_3\}),$$

for otherwise the edge of $S(y_1, a)$ incident on a is in T (since it is not in $A \cup B$), and we have a contradiction to the choice of a. Thus

$$a \in VS(y_1, w_1) \cup (VS(w_2, w_3) - \{w_2\}) \cup (VS(w_4, x_1) - \{w_4\}).$$

Now we distinguish three cases according to which set of this union contains a.

Suppose that $a \in VS(y_1, w_1)$. We already know that the trace of T_1 is a string over $\{0, 1, 2\}$. The symbols alternate between 2 and members of the set $\{0, 1\}$ for the trace of T_1 contains none of 00, 11, 22 by Lemma 13 and neither 01 nor 10 by Lemma 15. Therefore 2 is either the last symbol or the penultimate symbol in the trace of T_1. If 2 is the last symbol in the trace of T_1 we have a contradiction by Lemma 13, and if 2 is the penultimate symbol in the trace of T_1 then the last symbol of the trace of T_1 is either 0 or 1 and we have a contradiction by Lemma 10(a) and Lemma 17.

Suppose that $a \in VS(w_4, x_1) - \{w_4\}$. Note that a cannot be adjacent to w_4 since both vertices have indegree more than 1. Thus $b \in VS(w_4, a) - \{w_4\}$, for otherwise there would be a vertex of degree 2 in G. Therefore $\{a, b\} \subseteq VS(w_4, x_1)$ in contradition to Lemma 24.

Therefore $a \in VS(w_2, w_3) - \{w_2\}$. Then $b \in VS(w_2, a)$, for otherwise there would be a vertex of degree 2 in G. First we consider the case that the trace of S is 0. We define

$$T' = T_1 \cup S(a, x_1) \cup A(x_1, b) \cup T_2.$$

This is a directed path from y_2 to x_2 with

$$T' \cap (S(y_1, a) - (A \cup B)) = \emptyset.$$

By Theorem 5 we have

$$G = G[A \cup B \cup S' \cup T'],$$

a contradiction, since

$$S(w_1, w_2) \cap (A \cup B \cup S' \cup T') = \emptyset.$$

Now we consider the case that the trace of S is $0'$. By Lemma 13 and Lemma 15 the path $T_1(v, a)$ is an $\overline{A \cup B}$-arc. We define the following directed minimal path S'' from y_1 to x_1:

$$S'' = A(y_1, v) \cup T_1(v, a) \cup S(a, x_1).$$

The trace of S'' is $0'$. Let u be the origin of the $A(y_1, v_1)T_1$-arc with terminus v. Note that

$$T = T_1 \cup S(a, x_1) \cup A(x_1, y_1) \cup S(y_1, b) \cup T_2.$$

Then u is the first vertex in T that is also in S'' and w_1 is the last vertex in T that is also in S'', since the trace of T_2 does not contain 2. If we replace S by S'', this is a contradiction by Lemma 24 and finally shows that the trace of T_1 does not contain 2.

Now we know that the trace of T_1 is a string over $\{0, 1\}$. By Lemma 13 the trace of T_1 contains neither 00 nor 11, and by Lemma 15 it contains neither 01 nor 10. Therefore the trace of T_1 is either empty, 0 or 1. Since the directed path T_1 has property (A') the case that the trace of T_1 is 1 is not possible. ☐

Remark 2. *An argument similar to the one that showed that the trace of T_1 does not contain $0'$ leads to the following observation: If the trace of S is $0'$, the symbol 0 in the trace of T_1 corresponds to an $A(w_3, y_2)T_1$-arc and the symbol 0 in the trace of T_2 corresponds to an $A(x_2, w_2)T_2$-arc.*

Lemma 26. *Suppose $b <_S a$. Then $S(b, a) \subseteq S(w_2, w_3)$.*

Proof: The assertion can be deduced from Lemma 24 after we show that $S(b, a) \subseteq A \cup B$.

Define

$$X = S(y_1, b) \cup T_2$$

and

$$Y = T_1 \cup S(a, x_1).$$

Then X is a directed path from y_1 to x_2 and Y is a directed path from y_2 to x_1. Since $S(b,a) \cap (X \cup Y) = \emptyset$ and

$$G = G[A \cup B \cup X \cup Y]$$

by Theorem 5, we have $S(b,a) \subseteq A \cup B$. ☐

Lemma 27. *If the trace of* T_1 *is empty then* $a \notin VS(w_3, x_1)$.
If the trace of T_1 *is* 0 *then* $a \notin VS(y_1, w_3)$.
If the trace of T_2 *is empty then* $b \notin VS(y_1, w_2)$.
If the trace of T_2 *is* 0 *then* $b \notin VS(w_2, x_1)$.

Proof: By symmetry it suffices to show the assertions for T_1. The first assertion is an immediate consequence of Lemma 16.

Now suppose that the trace of T_1 is 0. If the trace of S is 0 then $a \notin VS(w_1, w_3)$ by Lemma 14; if the trace of S is $0'$ then $a \notin VS(w_1, w_3)$ by Lemma 13.

Suppose the trace of S is 0, and that $a \in VS(y_1, w_1)$. Let z_1, z_2, z_3 be the vertices of $VT_1 - \{y_2, a\}$ in the order in which they appear as T_1 is traced from y_2. Then

$$G[(A \cup B \cup S \cup T_1) - (A(y_2, z_1) \cup A(z_2, z_3) \cup B(x_2, y_1))]$$

is an even subdivision of $K_{3,3}$, in contradiction to the fact that G is minimal non-Pfaffian. If the trace of S is $0'$ then $a \notin VS(y_1, w_1)$ by Lemma 17 and Remark 2. ☐

The following lemma gives a complete list of graphs to be considered if $S \cap T \neq \emptyset$.

Lemma 28. *Suppose* $S \cap T \neq \emptyset$ *and that the directed path* T *has property* (A'). *Then one of the following cases is true:*

1. *the traces of* T_1 *and* T_2 *are* Λ *and* 0 *respectively,* $a \in VS(w_1, w_2) - \{w_2\}$, $b \in VS(w_1, w_2) - \{w_1\}$, $a <_S b$,
2. *the traces of* T_1 *and* T_2 *are empty,* $a \in VS(y_1, w_1)$, $b \in VS(w_2, w_3)$,
3. *the traces of* T_1 *and* T_2 *are empty,* $a \in VS(y_1, w_1)$, $b \in VS(w_4, x_1)$,
4. *the traces of* T_1 *and* T_2 *are empty,* $a \in VS(w_2, w_3)$, $b \in VS(w_2, w_3)$, $a <_S b$,
5. *the traces of* T_1 *and* T_2 *are empty,* $a \in VS(w_2, w_3)$, $b \in VS(w_2, w_3)$, $b <_S a$,
6. *the traces of* T_1 *and* T_2 *are empty,* $a \in VS(w_2, w_3)$, $b \in VS(w_4, x_1)$,
7. *the traces of* T_1 *and* T_2 *are* 0 *and* Λ *respectively,* $a \in VS(w_3, w_4) - \{w_4\}$, $b \in VS(w_3, w_4) - \{w_3\}$ *and* $a <_S b$.

Proof: First we deal with the case that

$$a \in (VS(w_1, w_2) - \{w_2\}) \cup (VS(w_3, w_4) - \{w_4\})$$

or

$$b \in (VS(w_1, w_2) - \{w_1\}) \cup (VS(w_3, w_4) - \{w_3\}).$$

Since there are no vertices of degree 2 in G, the vertex a is in $VS(w_1, w_2) - \{w_2\}$ if and only if $b \in VS(w_1, w_2) - \{w_1\}$. Furthermore $a <_S b$ in this case and by Lemma 27 the trace of T_1 is empty and the trace of T_2 is 0. This situation corresponds to the first case in the lemma.

Similarly $a \in VS(w_3, w_4) - \{w_4\}$ if and only if $b \in VS(w_3, w_4) - \{w_3\}$. In this case $a <_S b$, the trace of T_1 is 0 and the trace of T_2 is empty. This situation corresponds to the last case in the lemma.

Therefore we may now assume that

$$\{a,b\} \subset VS(y_1,w_1) \cup VS(w_2,w_3) \cup VS(w_4,x_1).$$

We show that the trace of T_1 is empty. Suppose the contrary, that the trace of T_1 is 0. By Lemma 27 and our assumption we have $a \in VS(w_4,x_1)$. By Lemma 26 we have $a <_S b$ and therefore $b \in VS(w_4,x_1)$. This is a contradiction to Lemma 24.

Similarly the trace of T_2 is empty.

By Lemma 27 and our assumption we have

$$a \in VS(y_1,w_1) \cup VS(w_2,w_3)$$

and

$$b \in VS(w_2,w_3) \cup VS(w_4,x_1).$$

From this result we deduce the following list of cases to be considered.

1. $a \in VS(y_1,w_1)$, $b \in VS(w_2,w_3)$
2. $a \in VS(y_1,w_1)$, $b \in VS(w_4,x_1)$
3. $a \in VS(w_2,w_3)$, $b \in VS(w_2,w_3)$, $a <_S b$
4. $a \in VS(w_2,w_3)$, $b \in VS(w_2,w_3)$, $b <_S a$
5. $a \in VS(w_2,w_3)$, $b \in VS(w_4,x_1)$.

□

Remark 3. *If we change from the reference orientation with respect to (M,N,f) to the reference orientation with respect to (N,M,f) the first case in Lemma 28 changes to the last case. Therefore we do not have to consider the last case. The same is true for the second case and the sixth case in Lemma 28, and so we do not consider the sixth case.*

5. The minimal non-Pfaffian near bipartite graphs

In this section we consider the cases in Lemma 22 and in Lemma 28, and with this complete the proof of Theorem 2. For that purpose we need the following lemma which has already been proved in [4].

Lemma 29. *Let G be a graph with a circuit C of odd length and let G^C be the graph obtained from G by contracting VC. If G^C is not Pfaffian, then neither is G.*

We divide the argument into cases according to whether or not $S \cap T = \emptyset$.

Case 1: In the case where $S \cap T = \emptyset$ it follows from Lemma 22 and Remark 1 that the trace of S may be assumed to be either 0 or $0'$ and that of T may be assumed to be one of 0, $0'$, 20, 21. Let the vertices in $VT - \{y_2,x_2\}$ be $z_1, z_2, \ldots, z_n$ in the order in which they appear.

Subcase 1.1: Suppose the trace of T is 0.

Subcase 1.1.1: Suppose the trace of S is 0. Consider the circuits

$$\begin{aligned} C_1 &= S(w_1,w_2) \cup A(w_2,w_1), \\ C_2 &= S(w_3,w_4) \cup A(w_4,w_3), \\ C_3 &= T(z_1,z_2) \cup A(z_2,z_1), \\ C_4 &= T(z_3,z_4) \cup A(z_4,z_3), \\ C_5 &= S \cup B(x_1,y_2) \cup T \cup B(x_2,y_1). \end{aligned}$$

Their sum is $A+B$. However, under our reference orientation all of C_1, C_2, C_3, C_4, C_5, $A+B$ are clockwise even, but under our extended Pfaffian orientation of H only $A+B$ is clockwise even, by Corollary 1. This result contradicts Lemma 12. *Subcase 1.1.2*: Suppose the trace of S is $0'$. By symmetry we may assume that $w_3 <_{A(x_2,y_2)} z_2$. Then G is isomorphic to Γ_1. The isomorphism ϕ from Γ_1 and G is given by $\phi(a)=x_2$, $\phi(b)=y_1$, $\phi(c)=w_1$, $\phi(d)=w_2$, $\phi(e)=w_3$, $\phi(f)=w_4$, $\phi(g)=x_1$, $\phi(h)=y_2$, $\phi(i)=z_1$, $\phi(j)=z_2$, $\phi(k)=z_3$, $\phi(l)=z_4$.

Subcase 1.2: Suppose the trace of T is $0'$.
Subcase 1.2.1: The case where the trace of S is 0 is symmetric to Subcase 1.1.2.
Subcase 1.2.2: Suppose the trace of S is $0'$. Consider the circuits

$$\begin{aligned} C_1 &= S \cup A(x_1,y_1),\\ C_2 &= T \cup A(x_2,y_2),\\ C_3 &= S \cup B(x_1,y_2) \cup T \cup B(x_2,y_1). \end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1.

Subcase 1.3: Suppose the trace of T is 20.
Subcase 1.3.1: Suppose the trace of S is 0. This case is similar to Subcase 1.1.1 except that vertices z_2, z_3, z_4 in Subcase 1.1.1 are replaced by z_4, z_5, z_6 respectively.
Subcase 1.3.2: Suppose the trace of S is $0'$. Note that $z_1=v_2$ and $z_3=v_1$. Contract the circuit $T(z_1,z_3)\cup\{e_2\}$. The resulting graph is isomorphic to Γ_1 whether or not $w_3 <_{A(x_2,y_2)} z_4$.

Subcase 1.4: Suppose the trace of T is 21.
Subcase 1.4.1: Suppose the trace of S is 0. Consider the circuits

$$\begin{aligned} C_1 &= S(w_1,w_2) \cup A(w_2,w_1),\\ C_2 &= S(w_3,w_4) \cup A(w_4,w_3),\\ C_3 &= T \cup A(x_2,y_2),\\ C_4 &= T(y_2,z_5) \cup B(z_5,y_2),\\ C_5 &= S \cup B(x_1,z_5) \cup T(z_5,x_2) \cup B(x_2,y_1). \end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1.
Subcase 1.4.2: Suppose the trace of S is $0'$. Contract the circuit $T(z_1,z_3)\cup\{e_2\}$. The resulting graph is isomorphic to Γ_1.

Case 2: If $S\cap T\neq\emptyset$ then by Remark 3 we see that only cases 1–5 in Lemma 28 need to be considered. Case i of the lemma is dealt with in Subcase 2.i below. Let the vertices in $(VT_1\cup VT_2)-\{y_2,x_2,a,b\}$ be $z_1,z_2,\ldots,z_n$ in the order in which they appear in T.
Subcase 2.1: The traces of T_1 and T_2 are Λ and 0 respectively, $a\in VS(w_1,w_2)-\{w_2\}$, $b\in VS(w_1,w_2)-\{w_1\}$, $a<_S b$.
Subcase 2.1.1: Suppose the trace of S is 0. Note that $z_1=v_2$ and $w_1=v_1$. Contraction of the circuit $T(z_1,a)\cup S(w_1,a)\cup\{e_2\}$ yields a graph isomorphic to Γ_1.
Subcase 2.1.2: Suppose the trace of S is $0'$. By Remark 2 we have $z_3 <_{A(x_2,y_2)} w_2$. Define

$$S' = S(y_1,b)\cup T(b,z_3)\cup A(z_3,w_2)\cup S(w_2,x_1).$$

By Theorem 5 we find that $G = G[A \cup B \cup S' \cup T]$, in contradiction to the fact that

$$S(b, w_2) \cap (A \cup B \cup S' \cup T) = \emptyset.$$

Subcase 2.2: The traces of T_1 and T_2 are empty, $a \in VS(y_1, w_1)$, $b \in VS(w_2, w_3)$. Note that $a = y_1$ and $b = w_3$.
Subcase 2.2.1: Suppose the trace of S is 0. Consider the circuits

$$\begin{aligned}
C_1 &= S(w_1, w_2) \cup A(w_2, w_1),\\
C_2 &= S(w_3, w_4) \cup A(w_4, w_3),\\
C_3 &= T_1 \cup S(y_1, w_3) \cup T_2 \cup A(x_2, y_2),\\
C_4 &= T_1 \cup S \cup B(x_1, y_2),\\
C_5 &= S(y_1, w_3) \cup T_2 \cup B(x_2, y_1).
\end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1.
Subcase 2.2.2: Suppose the trace of S is $0'$. Consider the circuits

$$\begin{aligned}
C_1 &= S \cup A(x_1, y_1),\\
C_2 &= T_1 \cup S(y_1, w_3) \cup A(w_3, y_2),\\
C_3 &= T_2 \cup A(x_2, w_3),\\
C_4 &= T_1 \cup S \cup B(x_1, y_2),\\
C_5 &= S(y_1, w_3) \cup T_2 \cup B(x_2, y_1).
\end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1.
Subcase 2.3: The traces of T_1 and T_2 are empty, $a \in VS(y_1, w_1)$, $b \in VS(w_4, x_1)$. Note that $a = y_1$ and $b = x_1$.
Subcase 2.3.1: Suppose the trace of S is 0. Consider the circuits

$$\begin{aligned}
C_1 &= S(w_1, w_2) \cup A(w_2, w_1),\\
C_2 &= S(w_3, w_4) \cup A(w_4, w_3),\\
C_3 &= T_1 \cup S \cup T_2 \cup A(x_2, y_2),\\
C_4 &= T_1 \cup S \cup B(x_1, y_2),\\
C_5 &= S \cup T_2 \cup B(x_2, y_1).
\end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1.
Subcase 2.3.2: Suppose the trace of S is $0'$. Consider the circuits

$$\begin{aligned}
C_1 &= S \cup A(x_1, y_1),\\
C_2 &= T_1 \cup S(y_1, w_3) \cup A(w_3, y_2),\\
C_3 &= S(w_2, x_1) \cup T_2 \cup A(x_2, w_2),\\
C_4 &= T_1 \cup S \cup B(x_1, y_2),\\
C_5 &= S \cup T_2 \cup B(x_2, y_1).
\end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1.
Subcase 2.4: The traces of T_1 and T_2 are empty, $a \in VS(w_2, w_3)$, $b \in VS(w_2, w_3)$, $a <_S b$. Note that $a = w_2$ and $b = w_3$.

Subcase 2.4.1: Suppose the trace of S is 0. Consider the circuits

$$\begin{aligned} C_1 &= S(w_1,w_2) \cup A(w_2,w_1), \\ C_2 &= S(w_3,w_4) \cup A(w_4,w_3), \\ C_3 &= T_1 \cup S(w_2,w_3) \cup T_2 \cup A(x_2,y_2), \\ C_4 &= T_1 \cup S(w_2,x_1) \cup B(x_1,y_2), \\ C_5 &= S(y_1,w_3) \cup T_2 \cup B(x_2,y_1). \end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1.

Subcase 2.4.2: Suppose the trace of S is $0'$. This case is symmetric to Subcase 2.4.1.

Subcase 2.5: The traces of T_1 and T_2 are empty, $a \in VS(w_2,w_3)$, $b \in VS(w_2,w_3)$, $b <_S a$.

Subcase 2.5.1: Suppose the trace of S is 0. Then G is isomorphic to Γ_2.

Subcase 2.5.2: Suppose the trace of S is $0'$. Consider the circuits

$$\begin{aligned} C_1 &= S \cup A(x_1,y_1), \\ C_2 &= T_1 \cup A(a,y_2), \\ C_3 &= T_2 \cup A(x_2,b), \\ C_4 &= T_1 \cup S(a,x_1) \cup B(x_1,y_2), \\ C_5 &= S(y_1,b) \cup T_2 \cup B(x_2,y_1). \end{aligned}$$

We obtain a contradiction by the method in Subcase 1.1.1. The proof of Theorem 2 is now complete.

6. NON-REDUCTION OF Γ_1 AND Γ_2 TO $K_{3,3}$

We conclude the paper by showing that neither Γ_1 nor Γ_2 is reducible to an even subdivision of $K_{3,3}$.

Lemma 30. *Both Γ_1 and Γ_2 are minimal non-Pfaffian graphs.*

Proof: Let Γ_1 and Γ_2 be oriented as in Figure 17.

First we consider Γ_1, which we have already seen to be non-Pfaffian. Suppose therefore that Γ_1 is not minimal. Let x be an edge such that $\Gamma_1 - \{x\}$ is non-Pfaffian. The 1-factors of Γ_1 are

$$\begin{aligned} f_1 &= \{(a,b),(c,d),(f,e),(h,g),(j,i),(l,k)\}, \\ f_2 &= \{(l,a),(b,c),(d,e),(g,f),(i,h),(k,j)\}, \\ f_3 &= \{(b,c),(a,d),(f,e),(h,g),(j,i),(l,k)\}, \\ f_4 &= \{(a,b),(c,d),(j,e),(g,f),(i,h),(l,k)\}, \\ f_5 &= \{(a,d),(b,c),(j,e),(g,f),(i,h),(l,k)\}, \\ f_6 &= \{(l,a),(b,c),(d,e),(g,f),(k,h),(j,i)\}, \\ f_7 &= \{(l,a),(b,g),(i,h),(k,j),(c,d),(f,e)\}, \\ f_8 &= \{(l,a),(b,g),(k,h),(c,d),(f,e),(j,i)\}, \\ f_9 &= \{(l,f),(d,e),(c,i),(k,j),(a,b),(h,g)\}, \\ f_{10} &= \{(a,d),(b,g),(c,i),(j,e),(k,h),(l,f)\}. \end{aligned}$$

Observe that the figure for the undirected graph Γ_1 is symmetric about the edge (l, f). Therefore we can assume that $x \notin f_1$. All 1-factors are associated with a plus sign except f_{10}. Thus $x \notin f_{10}$, for otherwise $\Gamma_1 - \{x\}$ is Pfaffian, and therefore $x \in f_2$.

Suppose that $x = (d, e)$ or $x = (k, j)$. We obtain a Pfaffian orientation of $\Gamma_1 - \{x\}$ if we change the orientation of (l, f), since every 1-factor of G that contains (l, f) also contains x except for f_{10}. If $x = (g, f)$ or $x = (i, h)$, we obtain a Pfaffian orientation of $\Gamma_1 - \{x\}$ by changing the orientation of (j, e). If $x = (l, a)$, we obtain a Pfaffian orientation of $\Gamma_1 - \{x\}$ by changing the orientation of (b, g). If $x = (b, c)$, we obtain a Pfaffian orientation of $\Gamma_1 - \{x\}$ by changing the orientation of (a, d). In all cases we have a contradiction to the fact that $\Gamma_1 - \{x\}$ was non-Pfaffian. Therefore Γ_1 is minimal non-Pfaffian.

Now suppose that the non-Pfaffian graph Γ_2 is not minimal, and let x be an edge such that $\Gamma_2 - \{x\}$ is non-Pfaffian. The 1-factors of Γ_2 are

$$\begin{aligned}
f_1 &= \{(a,b),(c,d),(e,f),(g,h),(i,j),(k,l)\},\\
f_2 &= \{(c,b),(e,d),(g,f),(i,h),(k,j),(a,l)\},\\
f_3 &= \{(a,b),(c,j),(e,d),(g,f),(i,h),(k,l)\},\\
f_4 &= \{(a,b),(c,d),(e,l),(g,f),(i,h),(k,j)\},\\
f_5 &= \{(c,b),(d,h),(k,g),(a,l),(e,f),(i,j)\},\\
f_6 &= \{(b,f),(e,d),(c,j),(a,i),(k,l),(g,h)\},\\
f_7 &= \{(b,f),(a,i),(k,j),(e,l),(c,d),(g,h)\},\\
f_8 &= \{(e,d),(b,f),(a,l),(k,g),(i,h),(c,j)\},\\
f_9 &= \{(g,f),(d,h),(c,b),(a,i),(k,j),(e,l)\},\\
f_{10} &= \{(b,f),(e,l),(k,g),(d,h),(c,j),(a,i)\}.
\end{aligned}$$

Observe that there is an automorphism of Γ_2 that interchanges the f_1-alternating circuits $f_1 + f_2$ and $f_1 + f_{10}$. Therefore we can assume that $x \in f_1 + f_{10}$. The figure for the undirected graph Γ_2 is symmetric about the line through the midpoints of the edges (a, b) and (g, h). Therefore we can assume that

$$\begin{gathered}
x \in (f_1 + f_{10}) - \{(b,f),(k,g),(c,j)\} = \\
\{(a,b),(c,d),(e,f),(g,h),(i,j),(k,l),(e,l),(d,h),(a,i)\}.
\end{gathered}$$

If $x \in f_1$ then the given orientation is a Pfaffian orientation of $\Gamma_2 - \{x\}$, since the sign of f_1 is the opposite of that of the other 1-factors. Therefore $x \in \{(e,l),(d,h),(a,i)\}$. If $x = (e, l)$, we obtain a Pfaffian orientation of $\Gamma_2 - \{x\}$ by changing the orientation of (c, d), since every 1-factor that contains (c, d) also contains x except for f_1. If $x = (d, h)$, we obtain a Pfaffian orientation of $\Gamma_2 - \{x\}$ by changing the orientation of (e, f). If $x = (a, i)$, we obtain a Pfaffian orientation of $\Gamma_2 - \{x\}$ by changing the orientation of (g, h). In all cases we have a contradiction to the fact that $\Gamma_2 - \{x\}$ was Pfaffian. Therefore Γ_2 is minimal non-Pfaffian.

Corollary 3. *Neither* Γ_1 *nor* Γ_2 *contains an even subdivision of* $K_{3,3}$.

Proof: If Γ_1 or Γ_2 contained an even subdivision of $K_{3,3}$ then Γ_1 or Γ_2 itself would be an even subdivision of $K_{3,3}$, since Γ_1 and Γ_2 are minimal non-Pfaffian and every even



FIGURE 17.

subdivision of $K_{3,3}$ is non-Pfaffian. But Γ_1 and Γ_2 both have 12 vertices of degree 3 whereas an even subdivision of $K_{3,3}$ has only 6. ☐

In order to see that Γ_1 and Γ_2 are not reducible to an even subdivision of $K_{3,3}$, we need the following two lemmas.

Lemma 31. *Let G be a minimal non-Pfaffian graph and C a circuit of odd length in G. Suppose that the graph G^C obtained by contracting VC to a vertex v is also non-Pfaffian. Let $VC = \{v_1, v_2, \ldots, v_n\}$. Then*

$$\deg v = \sum_{i=1}^{n} \deg v_i - 2n.$$

Moreover if w is a vertex in $VG - VC$ then w is also a vertex of G^C, and

$$\deg_G w = \deg_{G^C} w.$$

Proof: Let $N(u)$ denote the set of vertices in G that are adjacent to the vertex u in G. Moreover we assume that v_i is adjacent to v_{i+1} in C, for all $i < n$. We define $v_0 = v_n$ and $v_{n+1} = v_1$.

First we observe the following. Let $i, j \in \{1, 2, \ldots, n\}$ and $i \neq j$. We claim that

$$N(v_i) \cap N(v_j) \subseteq \{v_{i-1}, v_{i+1}\}.$$

Indeed, suppose $u \in N(v_i) \cap N(v_j)$. Then there are edges e_i and e_j joining u to v_i and v_j respectively. If $u \notin VC$, then the graph obtained from $G - \{e_i\}$ by contracting VC is G^C. Since G^C is supposed to be non-Pfaffian, it follows from Lemma 29 that $G - \{e_i\}$ is non-Pfaffian too. This is a contradiction to the fact that G was minimal non-Pfaffian. Therefore $u \in VC$. If $e_i \notin C$ or $e_j \notin C$ we can conclude in a similar way that either $G - \{e_i\}$ or $G - \{e_j\}$ is non-Pfaffian, and therefore have a contradiction again. Thus $\{e_i, e_j\} \subseteq C$, and the claim is proved. We infer that C has no chords, and any vertex not in C has at most one neighbour in C. We have

$$N(v) = \bigcup_{i=1}^{n} N(v_i) - VC = \bigcup_{i=1}^{n} (N(v_i) - \{v_{i-1}, v_{i+1}\}),$$

and so

$$\deg v = |N(v)| = \sum_{i=1}^{n} |N(v_i) - \{v_{i-1}, v_{i+1}\}| = \sum_{i=1}^{n} (|N(v_i)| - 2) = \sum_{i=1}^{n} \deg v_i - 2n.$$

Finally the degree of a vertex w that is not in C does not change upon contraction of C, since w is adjacent to at most one vertex in C. ☐

Lemma 32. *Let G be a minimal non-Pfaffian graph that is cubic and does not contain a circuit of length* 3*. Then G is not reducible to an even subdivision of $K_{3,3}$.*

Proof: Suppose the contrary, that is that there exists a sequence $C_0, C_1, \ldots, C_{p-1}$ of circuits of odd length and a sequence $G_0, G_1, \ldots, G_p$ of graphs such that $G_0 = G$, G_p is an even subdivision of $K_{3,3}$ and, for all $i < p$, G_{i+1} is obtained from G_i by contracting VC_i. We see inductively that for all i the graph G_i is minimal non-Pfaffian by Lemma 29, since G_p is non-Pfaffian and G_0 is minimal non-Pfaffian.

First we show that G_1 contains a vertex of degree at least 5. Since G_0 does not contain a circuit of length 3, the circuit C_0 must have length at least 5. Let $VC_0 = \{v_1, v_2, \ldots, v_n\}$ and let v be the corresponding vertex in G_1. Then, by Lemma 31,

$$\deg v = \sum_{i=1}^{n} \deg v_i - 2n = 3n - 2n = n \geq 5.$$

Again by Lemma 31 all the other vertices in G_1 have degree 3, since G_0 is cubic.

Now we show by induction that all the vertices in G_i, where $i \geq 1$, are of degree at least 3 and that there exists a vertex in G_i with degree at least 5. Therefore let us assume that the induction hypothesis is true for G_i and show it for G_{i+1}. First we show that every vertex in G_{i+1} is of degree at least 3. For all vertices in G_{i+1} except the one that corresponds to C_i this is an immediate consequence of Lemma 31 and the induction hypothesis. Let w be the vertex in G_{i+1} that corresponds to C_i and let $VC_i = \{w_1, w_2, \ldots, w_m\}$. Then, by Lemma 31 and the induction hypothesis,

$$\deg w = \sum_{i=1}^{m} \deg w_i - 2m \geq 3m - 2m = m \geq 3.$$

Now let u be the vertex of degree at least 5 in G_i. If $u \notin VC_i$ then $u \in VG_{i+1}$ and $\deg_{G_{i+1}} u = \deg_{G_i} u$, by Lemma 31. Therefore suppose that $u \in VC_i$, and without loss of generality assume that $u = w_1$. Then, by Lemma 31 and the induction hypothesis,

$$\deg w = \deg u + \sum_{i=2}^{m} \deg w_i - 2m \geq \deg u + 3(m-1) - 2m = \deg u + m - 3 \geq \deg u.$$

Therefore G_p contains a vertex of degree at least 5. This is a contradiction, since G_p was an even subdivision of $K_{3,3}$. ☐

Corollary 4. *Neither Γ_1 nor Γ_2 is reducible to an even subdivision of $K_{3,3}$.*

Proof: This is an immediate consequence of Lemma 32 since both Γ_1 and Γ_2 are minimal non-Pfaffian, cubic and do not contain a circuit of length 3. □

References

[1] P.W. Kasteleyn, Graph theory and crystal physics, in F. Harary, ed., "Graph Theory and Theoretical Physics", pp. 43-110, Acad. Press, New York, 1967.

[2] C.H.C. Little, Kasteleyn's theorem and arbitrary graphs, *Canad. J. Math.* **25** (1973), 758-764.

[3] C.H.C. Little, A characterization of convertible $(0,1)$-matrices, *J. Comb. Theory Ser. B* **18** (1975), 187-208.

[4] C.H.C. Little, F. Rendl and I. Fischer, Towards a Characterisation of Pfaffian graphs, preprint, `math.CO/9909026.`

[5] L. Lovász and M.D. Plummer, Matching Theory, Akadémiai Kiadó, Budapest, 1986.

[6] N. Robertson, P.D. Seymour and R. Thomas, Permanents, Pfaffian orientations, and even directed circuits, 1999, to appear in *Ann. Math.*